arXiv:0811.0926v3 [math.RT] 8 Mar 2010

On iterated almost ν-stable derived equivalences

Wei Hu

Abstract

In a recent paper [6], we introduced a classes of derived equivalences called almost ν-stable derived equivalences. The crucial property is that an almost ν-stable derived equivalence always induces a stable equivalence of Morita type, which generalizes a well-known result of Rickard: derived-equivalent self-injective algebras are stably equivalent of Morita type. In this paper, we shall consider the compositions of almost ν-stable derived equivalences and their quasi-inverses, which are called iterated almost ν-stable derived equivalences. We give a sufficient and necessary condition for a derived equivalence to be an iterated almost ν-stable derived equivalence, and give an explicit construction of the stable equivalence functor induced by an iterated almost ν-stable derived equivalence. As a consequence, we get some new sufficient conditions for a derived equivalence between general finite-dimensional algebras to induce a stable equivalence of Morita type.

1 Introduction

In [6], we introduced a class of derived equivalences called almost ν-stable derived equivalences. The crucial property is that an almost ν-stable derived equivalence always induces a stable equivalence of Morita type, which generalizes a classical result of Rickard ([13, Corollary 5.5]). This also gives a sufficient condition for a derived equivalence between general finite-dimensional algebras to induce a stable equivalence of Morita type. Note that many homological dimensions, such as global dimension, finitistic dimension, and representation dimension, are not invariant under derived equivalences in general. But they are all preserved by stable equivalences of Morita type. So, this also helps us to compare the homological dimensions of derived-equivalent algebras.

Let us first recall the definition of almost ν-stable derived equivalences. Let $F:\mathscr{D}^{\rm b}(A)\longrightarrow\mathscr{D}^{\rm b}(B)$ be a derived equivalence between two Artin algebras A and B, where $\mathscr{D}^{\rm b}(A)$ and $\mathscr{D}^{\rm b}(B)$ stand for the derived categories of bounded complexes over A and B, respectively. We use F^{-1} to denote a quasi-inverse of F. F is called an *almost* ν-*stable derived equivalence* if the following hold:

(1) *The tilting complex* $T^{\bullet}$ *associated to* F *has the following form:*

$$0\longrightarrow T^{-n}\longrightarrow\cdots\longrightarrow T^{-1}\longrightarrow T^{0}\longrightarrow 0$$

In this case, the tilting $\bar{T}^{\bullet}$ associated to F^{-1} has the following form (see [6, Lemma 2.1]):

$$0\longrightarrow \bar{T}^{0}\longrightarrow \bar{T}^{1}\longrightarrow\cdots\longrightarrow \bar{T}^{n}\longrightarrow 0$$

(2) $\mathrm{add}(\bigoplus_{i=1}^{n}T^{-i})=\mathrm{add}(\bigoplus_{i=1}^{n}\nu_A T^{-i})$ *and* $\mathrm{add}(\bigoplus_{i=1}^{n}\bar{T}^{i})=\mathrm{add}(\bigoplus_{i=1}^{n}\nu_B\bar{T}^{i})$, *where* ν *is the Nakayama functor.*

Let us remark that the composition of two almost ν-stable derived equivalences (or their quasi-inverses) is no longer almost ν-stable in general. If a derived equivalence is a composition $F\simeq F_1F_2\cdots F_m$ with F_i or F_i^{-1} being

---

2000 Mathematics Subject Classification: 18E30,16G10;18G20,16D90.

Keywords: derived equivalence, stable equivalence of Morita type, tilting complex.

an almost ν-stable derived equivalence for all i, then F is called an *iterated almost* ν*-stable derived equivalence*. By definition, we see that an almost ν-stable derived equivalence and its quasi-inverse are iterated almost ν-stable derived equivalences, and that the composition of two iterated almost ν-stable derived equivalences is again an iterated almost ν-stable derived equivalence. Clearly, an iterated almost ν-stable derived equivalence always induces a stable equivalence of Morita type, and therefore the involved algebras have many common homological dimensions. But the problem is:

Question: *Given a derived equivalence F, how to determine whether F is iterated almost* ν*-stable or not?*

The main purpose of this note is to give a complete answer to the above question. For a bounded complex $X^\bullet$ over an algebra A, we use $X^{\pm}$ to denote $\bigoplus_{i\neq 0} X^i$. The main result of this note can be stated as the following theorem.

Theorem 1.1. *Let $F : \mathscr{D}^{\rm b}(A) \longrightarrow \mathscr{D}^{\rm b}(B)$ be a derived equivalence between two Artin algebras A and B. Suppose that $T^\bullet$ and $\bar{T}^\bullet$ are the tilting complexes associated to F and F^{-1}, respectively. Then F is an iterated almost* ν*-stable derived equivalence if and only if* $\operatorname{add}({}_AT^{\pm}) = \operatorname{add}(\nu_A T^{\pm})$ *and* $\operatorname{add}({}_B\bar{T}^{\pm}) = \operatorname{add}(\nu_B\bar{T}^{\pm})$.

The above theorem tells us that, by checking the terms of tilting complexes, we can determine whether a derived equivalence is iterated almost ν-stable or not. If we work with finite-dimensional algebras over a field, then we have several other characterizations of iterated almost ν-stable derived equivalences. For details, see Theorem 3.6 below. As a consequence of Theorem 1.1, we have the following corollary, which provides a new sufficient condition for a derived equivalence to induce a stable equivalence of Morita type. For information on stable equivalences of Morita type, we refer to [3, 8, 9, 10].

Corollary 1.2. *Let $F : \mathscr{D}^{\rm b}(A) \longrightarrow \mathscr{D}^{\rm b}(B)$ be a derived equivalence between two finite-dimensional algebras A and B over a field. Suppose that $T^\bullet$ and $\bar{T}^\bullet$ are the tilting complexes associated to F and F^{-1}, respectively. If* $\operatorname{add}({}_AT^{\pm}) = \operatorname{add}(\nu_A T^{\pm})$ *and* $\operatorname{add}({}_B\bar{T}^{\pm}) = \operatorname{add}(\nu_B\bar{T}^{\pm})$, *then A and B are stably equivalent of Morita type.*

This paper is organized as follows. In Section 2, we shall recall some notations and basic facts. Theorem 1.1 will be proved in Section 3 after several lemmas. Section 4 is devoted to describing the stable equivalence functor induced by an iterated almost ν-stable derived equivalence. Finally, in Section 5, we shall give several methods to construct iterated almost ν-stable derived equivalences.

2 Preliminaries

In this section, we shall recall some basic definitions and facts needed in our later proofs.

Throughout this paper, unless specified otherwise, all algebras will be Artin algebras over a fixed commutative Artin ring R. All modules will be finitely generated unitary left modules. For an algebra A, the category of A-modules is denoted by A-mod; the full subcategory of A-mod consisting of projective modules is denoted by A-proj. The stable module category, denoted by A-$\underline{\rm mod}$, is the quotient category of A-mod modulo the ideal generated by morphisms factorizing through projective modules. We denote by ν_A the usual Nakayama functor.

Let $\mathcal{C}$ be an additive category. The composition of two morphisms $f : X \longrightarrow Y$ and $g : Y \longrightarrow Z$ in $\mathcal{C}$ will be denoted by fg. For two functors $F : \mathcal{C} \to \mathcal{D}$ and $G : \mathcal{D} \to \mathcal{E}$ of categories, their composition is denoted by GF. For an object X in $\mathcal{C}$, $\operatorname{add}(X)$ is the full subcategory of $\mathcal{C}$ consisting of all direct summands of finite direct sums of copies of X.

A complex $X^\bullet$ over $\mathcal{C}$ is a sequence $\cdots \longrightarrow X^{i-1} \xrightarrow{d_X^{i-1}} X^i \xrightarrow{d_X^i} X^{i+1} \xrightarrow{d_X^{i+1}} \cdots$ in $\mathcal{C}$ such that $d_X^i d_X^{i+1} = 0$ for all integers i. The category of complexes over $\mathcal{C}$ is denoted by $\mathscr{C}(\mathcal{C})$. The homotopy category of complexes over $\mathcal{C}$ is denoted by $\mathscr{K}(\mathcal{C})$. When $\mathcal{C}$ is an abelian category, the derived category of complexes over $\mathcal{C}$ is denoted by $\mathscr{D}(\mathcal{C})$. The full subcategory of $\mathscr{K}(\mathcal{C})$ and $\mathscr{D}(\mathcal{C})$ consisting of bounded complexes over $\mathcal{C}$ is denoted by $\mathscr{K}^{\rm b}(\mathcal{C})$

and $\mathscr{D}^{\mathrm{b}}(\mathcal{C})$, respectively. As usual, for a given algebra A, we simply write $\mathscr{K}^{\mathrm{b}}(A)$ and $\mathscr{D}^{\mathrm{b}}(A)$ for $\mathscr{K}^{\mathrm{b}}(A\text{-mod})$ and $\mathscr{D}^{\mathrm{b}}(A\text{-mod})$, respectively.

It is well-known that, for an algebra A, $\mathscr{K}^{\mathrm{b}}(A)$ and $\mathscr{D}^{\mathrm{b}}(A)$ are triangulated categories. For basic results on triangulated categories, we refer to Happel's book [4]. Throughout this paper, we use $X^{\bullet}[n]$ to denote the complex obtained by shifting $X^{\bullet}$ to the left by n degree.

Let A be an algebra. A homomorphism $f: X \longrightarrow Y$ of A-modules is called a *radical map* if, for any module Z and homomorphisms $h: Z \longrightarrow X$ and $g: Y \longrightarrow Z$, the composition hfg is not an isomorphism. A complex over A-mod is called a *radical* complex if all its differential maps are radical maps. Every complex over A-mod is isomorphic in the homotopy category $\mathscr{K}(A)$ to a radical complex. It is easy to see that if two radical complexes $X^{\bullet}$ and $Y^{\bullet}$ are isomorphic in $\mathscr{K}(A)$, then $X^{\bullet}$ and $Y^{\bullet}$ are isomorphic in $\mathscr{C}(A)$.

Two algebras A and B are said to be *derived-equivalent* if their derived categories $\mathscr{D}^{\mathrm{b}}(A)$ and $\mathscr{D}^{\mathrm{b}}(B)$ are equivalent as triangulated categories. In [12], Rickard proved that two algebras are derived-equivalent if and only if there is a complex $T^{\bullet}$ in $\mathscr{K}^{\mathrm{b}}(A\text{-proj})$ satisfying

(1) $\mathrm{Hom}(T^{\bullet}, T^{\bullet}[n]) = 0$ for all $n \neq 0$, and

(2) $\mathrm{add}(T^{\bullet})$ generates $\mathscr{K}^{\mathrm{b}}(A\text{-proj})$ as a triangulated category

such that $B \simeq \mathrm{End}(T^{\bullet})$. A complex in $\mathscr{K}^{\mathrm{b}}(A\text{-proj})$ satisfying the above two conditions is called a *tilting complex* over A. It is known that, given a derived equivalence F between A and B, there is a unique (up to isomorphism) tilting complex $T^{\bullet}$ over A such that $F(T^{\bullet}) \simeq B$. If $T^{\bullet}$ is a radical complex, it is called a tilting complex *associated* to F. Note that, by definition, a tilting complex associated to F is unique up to isomorphism in $\mathscr{C}^{b}(A)$.

The following lemma is useful in our later proof. For the convenience of the reader, we provide a proof.

Lemma 2.1. *Let $\mathcal{C}$ and $\mathcal{D}$ be two additive categories, and let $F: \mathscr{K}^{\mathrm{b}}(\mathcal{C}) \longrightarrow \mathscr{K}^{\mathrm{b}}(\mathcal{D})$ be a triangle functor. Let $X^{\bullet}$ be a complex in $\mathscr{K}^{\mathrm{b}}(\mathcal{C})$. For each term X^i, let $Y_i^{\bullet}$ be a complex isomorphic to $F(X^i)$. Then $F(X^{\bullet})$ is isomorphic to a complex $Z^{\bullet}$ with $Z^m = \bigoplus_{i+j=m} Y_i^j$ for all $m \in \mathbb{Z}$.*

Proof. We use induction on the number of non-zero terms of $X^{\bullet}$. If $X^{\bullet}$ has only one non-zero term, then it is obvious. Assume that $X^{\bullet}$ has more than one non-zero terms. Without loss of generality, we suppose that $X^{\bullet}$ is the following complex

$$0 \longrightarrow X^0 \longrightarrow X^1 \longrightarrow \cdots \longrightarrow X^n \longrightarrow 0$$

with $X^i \neq 0$ for all $i = 0, 1, \cdots, n$. Let $\sigma_{\geqslant 1} X^{\bullet}$ be the complex $0 \longrightarrow X^1 \longrightarrow \cdots \longrightarrow X^n \longrightarrow 0$. Then there is a distinguished triangle in $\mathscr{K}^{\mathrm{b}}(\mathcal{C})$:

$$X^0[-1] \longrightarrow \sigma_{\geqslant 1} X^{\bullet} \longrightarrow X^{\bullet} \longrightarrow X^0.$$

Applying F, we get a distinguished triangle in $\mathscr{K}^{\mathrm{b}}(\mathcal{D})$:

$$F(X^0[-1]) \longrightarrow F(\sigma_{\geqslant 1} X^{\bullet}) \longrightarrow F(X^{\bullet}) \longrightarrow F(X^0).$$

By induction, $F(\sigma_{\geqslant 1} X^{\bullet})$ is isomorphic to a complex $U^{\bullet}$ with $U^m = \bigoplus\limits_{1 \leqslant i \leqslant n, i+j=m} Y_i^j$. Thus, $F(X^{\bullet})$ is isomorphic to the mapping cone $Z^{\bullet}$ of the map from $Y_0^{\bullet}[-1]$ to $U^{\bullet}$. Thus, by definition, we have

$$Z^m = \bigoplus_{0 \leqslant i \leqslant n, i+j=m} Y_i^j = \bigoplus_{i+j=m} Y_i^j.$$

This finishes the proof. $\square$

Remark: Let $F: \mathscr{D}^{\mathrm{b}}(A) \longrightarrow \mathscr{D}^{\mathrm{b}}(B)$ be a derived equivalence between two algebras A and B. F induces an equivalence $F: \mathscr{K}^{\mathrm{b}}(A\text{-proj}) \longrightarrow \mathscr{K}^{\mathrm{b}}(B\text{-proj})$. So, for a bounded complex of projective A-modules, we can use the above lemma to calculate its image under F.

3 Characterizations of iterated almost ν-stable derived equivalences

In this section, we shall give a proof of our main result Theorem 1.1, which characterizes iterated almost ν-stable derived equivalences in terms of tilting complexes. In case that the algebras are finite-dimensional algebras over a field, we shall give several other characterizations of iterated almost ν-stable derived equivalences. For this purpose, we need some lemmas.

Let A be an algebra, and let ${}_AE$ be the direct sum of all those non-isomorphic indecomposable projective A-modules P with ν_A^iP being projective-injective for all $i \geqslant 0$. The A-module ${}_AE$ is unique up to isomorphism, and is called the *maximal* ν-*stable* A-module. If ${}_AQ$ is a projective A-module such that $\mathrm{add}({}_AQ) = \mathrm{add}(\nu_AQ)$, then clearly ${}_AQ \in \mathrm{add}({}_AE)$. Throughout this paper, we use ν_A-Stp to denote the category $\mathrm{add}({}_AE)$. Recall that for a bounded complex $X^\bullet$ over A, we use $X^\pm$ to denote the A-module $\bigoplus_{i\neq 0} X^i$.

Lemma 3.1. *Let $T^\bullet$ be a tilting complex associated to a derived equivalence $F : \mathscr{D}^{\mathrm{b}}(A) \longrightarrow \mathscr{D}^{\mathrm{b}}(B)$ between two algebras. Then the following two conditions are equivalent.*

(1) $\mathrm{add}(\nu_AT^\pm) = \mathrm{add}({}_AT^\pm)$;

(2) ${}_AT^\pm \in \nu_A$-Stp.

Proof. Clearly, we have (1) $\Rightarrow$ (2). It remains to show that (2) implies (1). Now we assume (2) holds. Let $Q_1 = \bigoplus_{i<0} T^i$. Using the same method in the proof of [6, Lemma 3.1], $F^{-1}(B)$ is isomorphic in $\mathscr{D}^{\mathrm{b}}(A)$ to a complex $X^\bullet$ with $X^i \in \mathrm{add}(\nu_AQ_1)$ for all $i < 0$. Thus, $T^\bullet \simeq X^\bullet$, and there is a quasi-isomorphism $f^\bullet : T^\bullet \longrightarrow X^\bullet$, which induces a quasi-isomorphism

$$\begin{tikzcd}
U^\bullet : & \cdots \arrow[r] & T^{-2} \arrow[r, "d_T^{-2}"] \arrow[d, "f^{-2}"] & T^{-1} \arrow[r, "\pi_T"] \arrow[d, "f^{-1}"] & \mathrm{Im}\, d_T^{-1} \arrow[r] \arrow[d, "f^0|_{\mathrm{Im}\, d_T^{-1}}"] & 0 \\
V^\bullet : & \cdots \arrow[r] & X^{-2} \arrow[r, "d_X^{-2}"] & X^{-1} \arrow[r, "\pi_X"] & \mathrm{Im}\, d_X^{-1} \arrow[r] & 0.
\end{tikzcd}$$

We claim that the canonical epimorphism $\pi_T : T^{-1} \longrightarrow \mathrm{Im}\, d_T^{-1}$ is still a radical map. Otherwise, let $h : Y \longrightarrow T^{-1}$ and $g : \mathrm{Im}\, d_T^{-1} \longrightarrow Y$ be such that $h\pi_Tg = 1_Y$. Then Y is isomorphic to a direct summand of T^{-1}, and therefore Y is an injective module. Thus, g factors through the inclusion $\lambda : \mathrm{Im}\, d_T^{-1} \longrightarrow T^0$, say $g = \lambda u$. Consequently $1_Y = h\pi_T\lambda u = hd_T^{-1}u$. This means that $d_T^{-1} : T^{-1} \longrightarrow T^0$ is not radical which is a contradiction. Since T^i and X^i are injective for all $i < 0$, by [6, Lemma 2.2], $U^\bullet$ and $V^\bullet$ are isomorphic in $\mathscr{K}^{\mathrm{b}}(A)$. Thus, T^i is a direct summand of X^i for all $i < 0$, and consequently $Q_1 = \bigoplus_{i<0} T^i \in \mathrm{add}(\nu_AQ_1)$. Since Q_1 and ν_AQ_1 have the same number of non-isomorphic indecomposable direct summands, we have $\mathrm{add}({}_AQ_1) = \mathrm{add}(\nu_AQ_1)$. Let $Q_2 := \bigoplus_{i>0} T^i$. Similarly, we have $\mathrm{add}({}_AQ_2) = \mathrm{add}(\nu_AQ_2)$. Consequently, $\mathrm{add}({}_AT^\pm) = \mathrm{add}({}_AQ_1 \oplus {}_AQ_2) = \mathrm{add}(\nu_AQ_1 \oplus \nu_AQ_2) = \mathrm{add}(\nu_AT^\pm)$. Hence (2) $\Rightarrow$ (1). $\square$

In the following, we shall use Lemma 3.1 freely. For instance, in the definition of an almost ν-stable equivalence, the condition $\mathrm{add}(\bigoplus_{i=1}^n T^{-i}) = \mathrm{add}(\bigoplus_{i=1}^n \nu_AT^{-i})$ is equivalent to say that $T^{-i} \in \nu_A$-Stp for all $i = 1, \cdots, n$.

Lemma 3.2. *Let $F : \mathscr{D}^{\mathrm{b}}(A) \longrightarrow \mathscr{D}^{\mathrm{b}}(B)$ be a derived equivalence between two algebras A and B, and let $T^\bullet$ and $\bar{T}^\bullet$ be the tilting complexes associated to F and F^{-1}, respectively. If* $\mathrm{add}({}_AT^\pm) = \mathrm{add}(\nu_AT^\pm)$ *and* $\mathrm{add}({}_B\bar{T}^\pm) = \mathrm{add}(\nu_B\bar{T}^\pm)$, *then F induces an equivalence between* $\mathscr{K}^{\mathrm{b}}(\nu_A\text{-Stp})$ *and* $\mathscr{K}^{\mathrm{b}}(\nu_B\text{-Stp})$.

Proof. Let ${}_AE$ (respectively, ${}_B\bar{E}$) be the maximal ν-stable A-module (respectively, B-module). Then by definition, we have $\nu_A\text{-Stp} = \mathrm{add}({}_AE)$ and $\nu_B\text{-Stp} = \mathrm{add}({}_B\bar{E})$. The complex $F({}_AE)$ is isomorphic to a complex $\bar{T}_1^\bullet$ in $\mathrm{add}(\bar{T}^\bullet)$. Since $\nu_AE \simeq {}_AE$, we have $\nu_B\bar{T}_1^\bullet \simeq \bar{T}_1^\bullet$ in $\mathscr{D}^{\mathrm{b}}(B)$. Hence there is a chain map η from $\bar{T}_1^\bullet$ to $\nu_B\bar{T}_1^\bullet$ such that the mapping cone $\mathrm{con}(\eta)$ is acyclic. By our assumption, all $\bar{T}_1^i$ and $\nu_B\bar{T}_1^i$ with $i \neq 0$ are projective-injective since they are all in ν_B-Stp. Hence $\mathrm{con}(\eta)$ splits, and therefore $\nu_B\bar{T}_1^0 \oplus \bar{Q}_1 \simeq \bar{T}_1^0 \oplus \bar{Q}_2$ for some $\bar{Q}_1, \bar{Q}_2 \in \nu_B$-Stp. Hence, $\nu_B\bar{T}_1^0 \in \mathrm{add}(\bar{T}_1^0 \oplus {}_B\bar{E})$. It follows that $\nu_B^i\bar{T}_1^0 \in \mathrm{add}(\bar{T}_1^0 \oplus {}_B\bar{E})$ is projective-injective for all $i \geqslant 0$. Hence

$\bar{T}_1^0 \in \nu_B$-Stp, and consequently $\bar{T}_1^\bullet$ is in $\mathscr{K}^{\rm b}(\nu_B\text{-Stp})$. Similarly, we can show that $F^{-1}({}_B\bar{E})$ is isomorphic to a complex in $\mathscr{K}^{\rm b}(\nu_A\text{-Stp})$ and the lemma is proved. □

The following lemma is useful in the proof of Theorem 1.1.

Lemma 3.3. *Let $F:\mathscr{D}^{\rm b}(A)\longrightarrow\mathscr{D}^{\rm b}(B)$ and $G:\mathscr{D}^{\rm b}(B)\longrightarrow\mathscr{D}^{\rm b}(C)$ be derived equivalences, and let $P^\bullet, \bar{P}^\bullet, Q^\bullet$, $\bar{Q}^\bullet$, $T^\bullet$, and $\bar{T}^\bullet$ be the tilting complexes associated to F, F^{-1}, G, G^{-1}, GF, and $F^{-1}G^{-1}$ respectively. If the following hold:*

(1) ${}_AP^{\pm}\in\nu_A$-Stp *and* ${}_B\bar{P}^{\pm}\in\nu_B$-Stp*;*

(2) ${}_BQ^{\pm}\in\nu_B$-Stp *and* ${}_C\bar{Q}^{\pm}\in\nu_C$-Stp*,*

then we have ${}_AT^{\pm}\in\nu_A$-Stp *and* ${}_C\bar{T}^{\pm}\in\nu_C$-Stp.

Proof. We only need to show that $\bar{T}^{\pm}\in\nu_C$-Stp, the other statement follows by symmetry. By definition, $\bar{T}^\bullet$ is isomorphic to $GF(A)\simeq G(\bar{P}^\bullet)$. Since $\bar{P}^i\in\nu_B$-Stp for all $i\neq 0$, by Lemma 3.2, $G(\bar{P}^i)$ is isomorphic to a complex $Y_i^\bullet$ in $\mathscr{K}^{\rm b}(\nu_C\text{-Stp})$ for all $i\neq 0$. For $i=0$, the complex $G(\bar{P}^0)$ is isomorphic to a complex $Y_0^\bullet$ in $\mathrm{add}(\bar{Q}^\bullet)$. By Lemma 2.1, the complex $G(\bar{P}^\bullet)$ is isomorphic to a complex $Z^\bullet$ with $Z^m=\bigoplus_{i+j=m}Y_i^j$. Since all Y_i^j, except Y_0^0, are in ν_C-Stp, we have $Z^{\pm}\in\nu_C$-Stp. Note that both $\bar{T}^\bullet$ and $Z^\bullet$ are in $\mathscr{K}^{\rm b}(C\text{-proj})$. The complexes $\bar{T}^\bullet$ and $Z^\bullet$ are isomorphic in $\mathscr{K}^{\rm b}(C\text{-proj})$. Furthermore, since the complex $\bar{T}^\bullet$ is a radical complex, it follows that $\bar{T}^i$ is a direct summand of Z^i for integers i, and consequently $\bar{T}^{\pm}\in\nu_C$-Stp. □

Finally, we have the following lemma which is crucial in the proof of our main result.

Lemma 3.4. *Let $F:\mathscr{D}^{\rm b}(A)\longrightarrow\mathscr{D}^{\rm b}(B)$ be a derived equivalence between two Artin algebras A and B, and let $T^\bullet$ be the associated tilting complex of F. If* ${}_AT^{\pm}\in\nu_A$-Stp*, then there is an almost ν-stable equivalence $G:\mathscr{D}^{\rm b}(C)\longrightarrow\mathscr{D}^{\rm b}(A)$ such that associated tilting complex $P^\bullet$ of FG satisfies that* $P^i\in\nu_C$-Stp *for all $i<0$ and $P^i=0$ for all $i>0$.*

Proof. Let ${}_AE$ be the maximal ν-stable A-module. Then $\nu_A\text{-Stp}=\mathrm{add}({}_AE)$. Suppose m is the maximal integer such that $T^m\neq 0$. By a dual statement of [5, Proposition 3.2], there is a tilting complex $Q^\bullet:=R^\bullet\oplus{}_AE[-m]$ over A, where $R^\bullet$ is of the form: $R^\bullet: 0\longrightarrow A\longrightarrow R^1\longrightarrow\cdots\longrightarrow R^m\longrightarrow 0$ with $R^i\in\nu_A$-Stp for all $i>0$. Let C be the endomorphism algebra of $Q^\bullet$, and let $H:\mathscr{D}^{\rm b}(A)\longrightarrow\mathscr{D}^{\rm b}(C)$ be a derived equivalence given by the tilting complex $Q^\bullet$. It is easy to see that $H({}_AE)\simeq{}_CP[m]$ for some ${}_CP\in\nu_C$-Stp, and $H(A)$ is isomorphic to a complex $S^\bullet$: $0\longrightarrow S^{-m}\longrightarrow\cdots\longrightarrow S^{-1}\longrightarrow S^0\longrightarrow 0$ with $S^i\in\nu_C$-Stp for all $i<0$. Let G is a quasi-inverse of H. Then $S^\bullet$ is a tilting complex associated to G. By Lemma 3.1, we see that G is almost ν-stable.

Now let $Y_i^\bullet:=H(T^i)$ for each integer i. Since $T^{\pm}\in\nu_A$-Stp, for each integer $i\neq 0$, we have $Y_i^\bullet\simeq P_i[m]$ for some $P_i\in\nu_C$-Stp. Moreover, $Y_i^\bullet=0$ for all $i>m$ since $T^i=0$ for all $i>m$. The complex $Y_0^\bullet$ has the property that $Y_0^i=0$ for all $i>0$ and $Y_0^i\in\nu_C$-Stp for all $i<0$. By Lemma 2.1, the complex $H(T^\bullet)$ is isomorphic to a complex $Z^\bullet$ with $Z^t=\bigoplus_{i+j=t}Y_i^j$. It follows that $Z^t=0$ for all $t>0$ and $Z^t\in\nu_C$-Stp for all $t<0$. Since $FG(H(T^\bullet))\simeq F(T^\bullet)\simeq B\simeq FG(P^\bullet)$ in $\mathscr{D}^{\rm b}(B)$, the complex $Z^\bullet$ is isomorphic in $\mathscr{D}^{\rm b}(C)$ to the tilting complex $P^\bullet$ associated to FG. Since both $Z^\bullet$ and $P^\bullet$ are in $\mathscr{K}^{\rm b}(C\text{-proj})$, they are isomorphic in $\mathscr{K}^{\rm b}(C\text{-proj})$. Since $P^\bullet$ is a radical complex, the term P^i is a direct summand of Z^i for all i, and consequently $P^\bullet$ has the desired property. □

We are now in the position to give a proof of our main result.

Proof of Theorem 1.1. Assume that F is an iterated almost ν-stable derived equivalence. Let $F\simeq F_1F_2\cdots F_m$ be a composition such that F_i or F_i^{-1} is an almost ν-stable derived equivalence. Then by Lemma 3.3, we have $\mathrm{add}({}_AT^{\pm})=\mathrm{add}(\nu_AT^{\pm})$ and $\mathrm{add}({}_B\bar{T}^{\pm})=\mathrm{add}(\nu_B\bar{T}^{\pm})$. Conversely, assume that $\mathrm{add}({}_AT^{\pm})=\mathrm{add}(\nu_AT^{\pm})$ and $\mathrm{add}({}_B\bar{T}^{\pm})=\mathrm{add}(\nu_B\bar{T}^{\pm})$. By Lemma 3.4, there is an almost ν-stable derived equivalence $G:\mathscr{D}^{\rm b}(C)\longrightarrow\mathscr{D}^{\rm b}(A)$ such that the tilting complex $P^\bullet$ associated to FG has the property that $P^i=0$ for all $i>0$ and $P^i\in\nu_C$-Stp for all $i<0$. By Lemma 3.1, we have $\mathrm{add}(\bigoplus_{i<0}P^i)=\mathrm{add}(\bigoplus_{i<0}\nu_CP^i)$. Let $\bar{P}^\bullet$ be the tilting complex associated to $G^{-1}F^{-1}$. By Lemma 3.3, we have $\mathrm{add}({}_B\bar{P}^{\pm}) = \mathrm{add}(\nu_B\bar{P}^{\pm})$. Since $P^i = 0$ for all $i > 0$, by [6, Lemma 2.1], we have $\bar{P}^i = 0$ for all $i < 0$. Hence FG is an almost ν-stable derived equivalence. Thus, $F \simeq (FG)G^{-1}$ is an iterated almost ν-stable derived equivalence. $\square$

Remark: (1) Theorem 1.1 gives us a method to determine whether a derived equivalence is iterated almost ν-stable or not by checking the terms of the involved tilting complexes.

(2) Let P be a projective A-module. The condition $\mathrm{add}({}_AP) = \mathrm{add}(\nu_AP)$ is equivalent to say that P is projective-injective and $\mathrm{add}(\mathrm{top}(P)) = \mathrm{add}(\mathrm{soc}(P))$.

(3) The proof of Corollary 1.2 follows immediately from Theorem 1.1 and [10, Theorem 5.3]. But Corollary 1.2 generalizes [10, Theorem 5.3]. This gives a new sufficient condition for a derived equivalence to induce a stable equivalence of Morita type.

As an application of Theorem 1.1, we have the following corollary.

Corollary 3.5. *Let $F : \mathscr{D}^{\mathrm{b}}(A) \longrightarrow \mathscr{D}^{\mathrm{b}}(B)$ be a derived equivalence between two Artin algebras A and B, and let $T^{\bullet}$ and $\bar{T}^{\bullet}$ be the tilting complexes associated to F and F^{-1}, respectively. If $\mathrm{add}({}_AT^{\pm}) = \mathrm{add}(\nu_AT^{\pm})$ and $\mathrm{add}({}_B\bar{T}^{\pm}) = \mathrm{add}(\nu_B\bar{T}^{\pm})$, then the following hold:*

(1) $\mathrm{fin.dim}(A) = \mathrm{fin.dim}(B)$, *and* $\mathrm{gl.dim}(A) = \mathrm{gl.dim}(B)$*;*

(2) $\mathrm{rep.dim}(A) = \mathrm{rep.dim}(B)$*;*

(3) $\mathrm{dom.dim}(A) = \mathrm{dom.dim}(B)$,

where $\mathrm{fin.dim}, \mathrm{gl.dim}, \mathrm{rep.dim}$ *and* $\mathrm{dom.dim}$ *stand for finitistic dimension, global dimension, representation dimension and dominant dimension, respectively.*

Proof. The corollary follows from [6, Corollary 1.2] and Theorem 1.1. $\square$

Now we work with finite-dimensional algebras over a field. In this case, we get several other characterizations of iterated almost ν-stable derived equivalences, which is the following theorem.

Theorem 3.6. *Let $F : \mathscr{D}^{\mathrm{b}}(A) \longrightarrow \mathscr{D}^{\mathrm{b}}(B)$ be a derived equivalence between two finite-dimensional basic algebras A and B over a field, and let $T^{\bullet}$ and $\bar{T}^{\bullet}$ be the tilting complexes associated to F and F^{-1}, respectively. Then the following are equivalent:*

(1) *The functor F is an iterated almost ν-stable derived equivalence.*

(2) $\mathrm{add}(\nu_AT^{\pm}) = \mathrm{add}({}_AT^{\pm})$ *and* $\mathrm{add}(\nu_B\bar{T}^{\pm}) = \mathrm{add}({}_B\bar{T}^{\pm})$.

(3) $T^{\pm} \in \nu_A\text{-}\mathrm{Stp}$ *and* $\bar{T}^{\pm} \in \nu_B\text{-}\mathrm{Stp}$.

(4) *For each indecomposable projective A-module $P \notin \nu_A\text{-}\mathrm{Stp}$, the image $F(\mathrm{top}(P))$ is isomorphic in $\mathscr{D}^{\mathrm{b}}(B)$ to a simple B-module.*

(5) *For each indecomposable projective A-module $P \notin \nu_A\text{-}\mathrm{Stp}$, the following conditions are satisfied:*

  (a) *$P \notin \mathrm{add}({}_AT^{\pm})$;*

  (b) *the multiplicity of P as a direct summand of ${}_AT^0$ is* 1.

Proof. It follows from Theorem 1.1 and Lemma 3.1 that the statements (1), (2) and (3) are equivalent. Note that for any simple module S over a basic algebra Λ, the dimension of S as an $\mathrm{End}_{\Lambda}(S)$-space is 1. In this proof, let ${}_AE$ and ${}_B\bar{E}$ be the maximal ν-stable A-module and B-module, respectively.

(4) $\Rightarrow$ (5) For each indecomposable projective A-module P not in ν_A-Stp, since $F(\mathrm{top}(P))$ is isomorphic in $\mathscr{D}^{\mathrm{b}}(B)$ to a simple B-module, we have $\mathrm{Hom}_{\mathscr{D}^{\mathrm{b}}(A)}(T^{\bullet}, \mathrm{top}(P)[i]) = 0$ for all $i \neq 0$, and

$$\mathrm{Hom}_{\mathscr{D}^{\mathrm{b}}(A)}(T^{\bullet}, \mathrm{top}(P)) \simeq \mathrm{Hom}_B(B, F(\mathrm{top}(P))) \simeq F(\mathrm{top}(P))$$

is one-dimensional over the division ring $\mathrm{End}_A(\mathrm{top}(P))$. Note that $T^{\bullet}$ is a radical complex. It follows that P is not a direct summand of $T^{\pm}$ and the multiplicity of P as a direct summand of T^0 is 1.

(5) $\Rightarrow$ (4) By condition (a), we see that $\mathrm{Hom}_{\mathscr{D}^{\mathrm{b}}(A)}(T^{\bullet},\mathrm{top}(P)[i])=0$ for all $i\neq 0$. Hence $F(\mathrm{top}(P))$ is isomorphic to an indecomposable B-module X. By condition (b), we can assume that $T_P^{\bullet}$ is the only indecomposable direct summand of $T^{\bullet}$ such that P is a direct summand of its degree zero term. Suppose that $\bar{P}$ is the indecomposable projective B-module corresponding to the direct summand $T_P^{\bullet}$. Then

$$\mathrm{Hom}_B(B,X)\simeq \mathrm{Hom}_{\mathscr{D}^{\mathrm{b}}(A)}(T^{\bullet},\mathrm{top}(P))\simeq \mathrm{Hom}_{\mathscr{D}^{\mathrm{b}}(A)}(T_P^{\bullet},\mathrm{top}(P))\simeq \mathrm{Hom}_B(\bar{P},X).$$

This implies that X only contains $\mathrm{top}(\bar{P})$ as composition factors. If X is not a simple B-module, then there is a nonzero map $X\longrightarrow \mathrm{soc}(X)\longrightarrow X$ in $\mathrm{End}_B(X)$ which is not an isomorphism. This contradicts to the fact that $\mathrm{End}_B(X)\simeq \mathrm{End}_A(\mathrm{top}(P))$ is a division ring. Hence $X\simeq F(\mathrm{top}(P))$ is a simple B-module.

(3) $\Rightarrow$ (4) By definition, we have $\mathrm{add}({}_AE)=\nu_A$-Stp and $\mathrm{add}({}_B\bar{E})=\nu_B$-Stp. Let P be an indecomposable projective A-module not in ν_A-Stp. Then it is clear that $\mathrm{Hom}_{\mathscr{D}^{\mathrm{b}}(A)}(T^{\bullet},\mathrm{top}(P)[i])=0$ for all $i\neq 0$ since $T^{\pm}\in\nu_A$-Stp, and consequently $F(\mathrm{top}(P))$ is isomorphic in $\mathscr{D}^{\mathrm{b}}(B)$ to a B-module X. By Lemma 3.2, the complex $F^{-1}({}_B\bar{E})$ is isomorphic in $\mathscr{D}^{\mathrm{b}}(A)$ to a complex $E^{\bullet}$ in $\mathscr{K}^{\mathrm{b}}(\nu_A\text{-Stp})$. Hence

$$\mathrm{Hom}_B({}_B\bar{E},X)\simeq \mathrm{Hom}_{\mathscr{D}^{\mathrm{b}}(A)}(F^{-1}({}_B\bar{E}),\mathrm{top}(P))\simeq \mathrm{Hom}_{\mathscr{K}^{\mathrm{b}}(A)}(E^{\bullet},\mathrm{top}(P))=0.$$

If ${}_BX$ is not simple, then there is a short exact sequence $0\longrightarrow\bar{U}\longrightarrow X\longrightarrow\bar{V}\longrightarrow 0$ in B-module with $\bar{U},\bar{V}$ non-zero. Applying $\mathrm{Hom}_B({}_B\bar{E},-)$, we get that $\mathrm{Hom}_B({}_B\bar{E},\bar{U})=0=\mathrm{Hom}_B({}_B\bar{E},\bar{V})$, and consequently $\mathrm{Hom}_{\mathscr{D}^{\mathrm{b}}(B)}(\bar{T}^{\bullet},\bar{U}[i])=0=\mathrm{Hom}_{\mathscr{D}^{\mathrm{b}}(B)}(\bar{T}^{\bullet},\bar{V}[i])$ for all $i\neq 0$ since $\bar{T}^{\pm}\in\nu_B$-Stp. Hence $F^{-1}(\bar{U})$ and $F^{-1}(\bar{V})$ are isomorphic to A-modules U and V, respectively. Thus, we get a distinguished triangle

$$U\longrightarrow \mathrm{top}(P)\longrightarrow V\longrightarrow U[1]$$

in $\mathscr{D}^{\mathrm{b}}(A)$ by applying F^{-1} to the distinguished triangle $\bar{U}\longrightarrow X\longrightarrow\bar{V}\longrightarrow\bar{U}[1]$. Applying $\mathrm{Hom}_{\mathscr{D}^{\mathrm{b}}(A)}(A,-)$ to the above triangle, we get an exact sequence $0\longrightarrow U\longrightarrow\mathrm{top}(P)\longrightarrow V\longrightarrow 0$ with non-zero A-modules U and V. This contradicts to the fact that $\mathrm{top}(P)$ is a simple A-module. Hence $F(\mathrm{top}(P))\simeq X$ is a simple B-module.

(4) $\Rightarrow$ (3) For each indecomposable projective A-module P not in ν_A-Stp, since $F(\mathrm{top}(P))$ is isomorphic $\mathscr{D}^{\mathrm{b}}(B)$ to a simple B-module, we have $\mathrm{Hom}_{\mathscr{D}^{\mathrm{b}}(A)}(T^{\bullet},\mathrm{top}(P)[i])=0$ for all $i\neq 0$. Together with the isomorphism

$$\mathrm{Hom}_{\mathscr{D}^{\mathrm{b}}(A)}(T^{\bullet},\mathrm{top}(P)[i])\simeq \mathrm{Hom}_{\mathscr{K}^{\mathrm{b}}(A)}(T^{\bullet},\mathrm{top}(P)[i])\simeq \mathrm{Hom}_A(T^i,\mathrm{top}(P)),$$

we get $\mathrm{Hom}_A(T^i,\mathrm{top}(P))=0$ for all $i\neq 0$ and for all indecomposable projective A-module P not in ν_A-Stp. Hence $T^i\in\nu_A$-Stp for all $i\neq 0$, that is, $T^{\pm}\in\nu_A$-Stp. Now let ${}_AQ$ be a projective A-module such that ${}_AA\simeq{}_AE\oplus{}_AQ$. It follows by assumption that $F(\mathrm{top}(Q))$ is a semi-simple B-module. Suppose that $\bar{Q}$ is a projective cover of $F(\mathrm{top}(Q))$, and suppose that ${}_BB\simeq\bar{Q}\oplus W$. Since $\bar{T}^{\bullet}$ is a radical complex in B-proj, we have

$$\begin{aligned}\mathrm{Hom}_B(\bar{T}^i,\mathrm{top}(\bar{Q})) &\simeq \mathrm{Hom}_{\mathscr{K}^{\mathrm{b}}(B)}(\bar{T}^{\bullet},\mathrm{top}(\bar{Q})[i])\\ &\simeq \mathrm{Hom}_{\mathscr{D}^{\mathrm{b}}(B)}(\bar{T}^{\bullet},\mathrm{top}(\bar{Q})[i])\\ &\simeq \mathrm{Hom}_{\mathscr{D}^{\mathrm{b}}(A)}(A,\mathrm{top}(Q)[i])=0\end{aligned}$$

for all $i\neq 0$. Hence $\bar{T}^{\pm}\in\mathrm{add}({}_BW)$. It remains to show ${}_BW\in\nu_B$-Stp. Note that $\mathrm{Hom}_B({}_BW,\mathrm{top}(\bar{Q})[i])=0$ for all integers i. It follows that $\mathrm{Hom}_{\mathscr{D}^{\mathrm{b}}(A)}(F^{-1}({}_BW),\mathrm{top}(Q)[i])=0$ for all integers i. Let $L^{\bullet}$ be a radical complex in $\mathscr{K}^{\mathrm{b}}(A\text{-proj})$ such that $F^{-1}({}_BW)\simeq L^{\bullet}$. Then $\mathrm{Hom}_A(L^i,\mathrm{top}(Q))\simeq\mathrm{Hom}_{\mathscr{D}^{\mathrm{b}}(A)}(L^{\bullet},\mathrm{top}(Q)[i])=0$ for all integers i. Hence $L^i\in\mathrm{add}({}_AE)$ for all integers i. Using the same proof as the proof of [1, Theorem 2.1], we can show that ν_B^iW is a projective B-module for all $i\geqslant 0$. It follows that ${}_BW\simeq\nu_B^kW$ for some $k>0$. Hence ${}_BW$ is projective-injective and ν_B^iW is projective-injective for $i>0$, and consequently ${}_BW\in\nu_B$-Stp. This finishes the proof. $\square$

Remark: (1) By Theorem 3.6 (5), we see that if we consider finite-dimensional algebras over a field, then we can determine whether a derived equivalence F is iterated almost ν-stable or not by checking the terms of the tilting complex associated to F, and we do not need to check the terms of the tilting complex associated to F^{-1}, which is needed in Theorem 1.1.

(2) It is interesting to know whether Theorem 3.6 holds for general Artin algebras. Note that the only problem is the step "(4) ⇒ (3)", where the method in the proof of [1, Theorem 2.1] does not work for general Artin algebras.

As a consequence, we have the following corollary.

Corollary 3.7. *Let $F: \mathscr{D}^{\mathrm{b}}(A) \longrightarrow \mathscr{D}^{\mathrm{b}}(B)$ be a derived equivalence between two finite-dimensional basic algebras over a field. If one of the equivalent conditions in* Theorem 3.6 *is satisfied, then the algebras A and B are stably equivalent of Morita type.*

Proof. This follows from Theorem 3.6, and [6, Theorem 5.3]. ☐

We end this section by using a simple example to illustrate Theorem 3.6 and Corollary 3.7.

Example: Let k be a field, and let A and B be finite-dimensional k-algebras given by quivers with relations in Fig. 1 and Fig. 2, respectively.



$$\alpha\beta\gamma = \beta\gamma\alpha\beta = \gamma\alpha\beta\gamma = 0$$

Fig. 1



$$\alpha\gamma = \delta\beta = 0$$
$$\alpha\beta = \delta\gamma\delta = \beta\alpha - \gamma\delta = 0.$$

Fig. 2

Let $P(i)$ denote the indecomposable projective A-module corresponding to the vertex i. Then there is a tilting complex of A-modules

$$T^\bullet: \quad 0 \longrightarrow P(2) \oplus P(2) \oplus P(3) \overset{[f,0,0]^T}{\longrightarrow} P(1) \longrightarrow 0$$

with $P(1)$ in degree zero. One can check that $\mathrm{End}_{\mathscr{K}^{\mathrm{b}}(A\text{-proj})}(T^\bullet)$ is isomorphic to B, and that $\nu_A\text{-Stp} = \mathrm{add}(P(2) \oplus P(3))$. Hence the tilting complex satisfies the condition (5) in Theorem 3.6. Therefore, the complex $T^\bullet$ induces an iterated almost ν-stable derived equivalence (actually even an almost ν-stable derived equivalence) between A and B. By Corollary 3.7, the algebras A and B are stably equivalent of Morita type.

4 The stable equivalence functor

In this section, we will give a description of the stable equivalence functor induced by an iterated almost ν-stable derived equivalence.

Let A be an Artin algebra, and let ${}_AE$ be a maximal ν-stable A-module. Then by definition $\nu_A\text{-Stp} = \mathrm{add}({}_AE)$. We use $A\text{-}\underline{\mathrm{mod}}^{\nu}$ to denote the quotient category of A-mod modulo morphisms factorizing through modules in ν_A-Stp. The Hom-space in $A\text{-}\underline{\mathrm{mod}}^{\nu}$ is denoted by $\underline{\mathrm{Hom}}^{\nu}_A(-,-)$. For a morphism f in A-mod, its image in $A\text{-}\underline{\mathrm{mod}}^{\nu}$ under the canonical functor from A-mod to $A\text{-}\underline{\mathrm{mod}}^{\nu}$ is denoted by $\underline{f}$. The category $\mathscr{K}^{\mathrm{b}}(\nu_A\text{-Stp})$ is a clearly thick subcategory (that is, a triangulated full subcategory closed under taking direct summands) of $\mathscr{D}^{\mathrm{b}}(A)$. Let $\mathscr{D}^{\mathrm{b}}(A)/\mathscr{K}^{\mathrm{b}}(\nu_A\text{-Stp})$ be the Verdier quotient category, then we have a canonical additive functor

$$\Sigma': A\text{-mod} \longrightarrow \mathscr{D}^{\mathrm{b}}(A)/\mathscr{K}^{\mathrm{b}}(\nu_A\text{-Stp})$$

obtained by composing the natural embedding from A-mod to $\mathscr{D}^{\mathrm{b}}(A)$ and the quotient functor from $\mathscr{D}^{\mathrm{b}}(A) \longrightarrow \mathscr{D}^{\mathrm{b}}(A)/\mathscr{K}^{\mathrm{b}}(\nu_A\text{-Stp})$. For the definition and basic properties of Verdier quotient, we refer to [11, Chapter 2].

Since $\Sigma'({}_AE)$ is clearly isomorphic to zero object in $\mathscr{D}^{\mathrm{b}}(A)/\mathscr{K}^{\mathrm{b}}(\nu_A\text{-Stp})$, the functor Σ' induces an additive functor

$$\Sigma : A\text{-}\underline{\mathrm{mod}}^{\nu} \longrightarrow \mathscr{D}^{\mathrm{b}}(A)/\mathscr{K}^{\mathrm{b}}(\nu_A\text{-Stp}).$$

Keeping this notation, we have a proposition, which can be viewed as a generalization of a well-known result of Rickard [14, Theorem 2.1]

Proposition 4.1. *The functor*

$$\Sigma : A\text{-}\underline{\mathrm{mod}}^{\nu} \longrightarrow \mathscr{D}^{\mathrm{b}}(A)/\mathscr{K}^{\mathrm{b}}(\nu_A\text{-Stp})$$

is fully faithful. Moreover, the functor Σ is an equivalence if and only if A is self-injective.

Proof. A morphism $X^\bullet \longrightarrow Y^\bullet$ in $\mathscr{D}^{\mathrm{b}}(A)/\mathscr{K}^{\mathrm{b}}(\nu_A\text{-Stp})$ is denoted by a fraction $s^{-1}a : X^\bullet \xLeftarrow{s} Z^\bullet \xrightarrow{a} Y^\bullet$, where a and s are morphisms in $\mathscr{D}^{\mathrm{b}}(A)$, and if $Z^\bullet \xRightarrow{s} X^\bullet \longrightarrow U^\bullet \longrightarrow Z^\bullet[1]$ is a distinguished triangle in $\mathscr{D}^{\mathrm{b}}(A)$, then $U^\bullet \in \mathscr{K}^{\mathrm{b}}(\nu_A\text{-Stp})$. A morphism s' in $\mathscr{D}^{\mathrm{b}}(A)$ with this property will be denoted by $\xRightarrow{s'}$. Two morphisms $X^\bullet \xLeftarrow{s} U^\bullet \xrightarrow{a} Y^\bullet$ and $X^\bullet \xLeftarrow{r} V^\bullet \xrightarrow{b} Y^\bullet$ are equal if and only if there are morphisms $W^\bullet \xRightarrow{t} U^\bullet$ and $W^\bullet \xRightarrow{h} V^\bullet$ such that $ts = hr$ and $ta = hb$. An isomorphism from X to Y is of the form $X \xLeftarrow{s} U^\bullet \xRightarrow{t} Y$.

First, we show that Σ is a full functor. For this purpose, it suffices to show that Σ' is a full functor. Now let $f : X \longrightarrow Y$ be a morphism in A-mod. Then $\Sigma'(f)$ is the morphism $X \xLeftarrow{1_X} X \xrightarrow{f} Y$. We need to show that each morphism from X to Y in $\mathscr{D}^{\mathrm{b}}(A)/\mathscr{K}^{\mathrm{b}}(\nu_A\text{-Stp})$ is of this form. Let $X \xLeftarrow{s} U^\bullet \xrightarrow{a} Y$ be a morphism in $\mathscr{D}^{\mathrm{b}}(A)/\mathscr{K}^{\mathrm{b}}(\nu_A\text{-Stp})$. By definition, there is a distinguished triangle $U^\bullet \xrightarrow{s} X \xrightarrow{g} E^\bullet \longrightarrow U^\bullet[1]$ in $\mathscr{D}^{\mathrm{b}}(A)$ with $E^\bullet \in \mathscr{K}^{\mathrm{b}}(\nu_A\text{-Stp})$. Consider the distinguished triangle in $\mathscr{D}^{\mathrm{b}}(A)$

$$\sigma_{\geqslant 0}E^\bullet \xrightarrow{\alpha} E^\bullet \xrightarrow{\beta} \sigma_{<0}E^\bullet \longrightarrow (\sigma_{\geqslant 0}E^\bullet)[1].$$

Since $E^\bullet$ is clearly in $\mathscr{K}^{\mathrm{b}}(A\text{-inj})$, we have $\mathrm{Hom}_{\mathscr{D}^{\mathrm{b}}(A)}(X, \sigma_{<0}E^\bullet) \simeq \mathrm{Hom}_{\mathscr{K}^{\mathrm{b}}(A)}(X, \sigma_{<0}E^\bullet) = 0$. It follows that $g\beta = 0$, and therefore g factorizes through α. Hence we can form the following commutative diagram in $\mathscr{D}^{\mathrm{b}}(A)$ with rows being distinguished triangles.

$$\begin{tikzcd}
V^\bullet \arrow[r, Rightarrow, "h"] \arrow[d, Rightarrow, "r"] & X \arrow[r] \arrow[d, equal] & \sigma_{\geqslant 0}E^\bullet \arrow[r, "w"] \arrow[d, "\alpha"] & V^\bullet[1] \arrow[d, Rightarrow, "r[1]"] \\
U^\bullet \arrow[r, Rightarrow, "s"] & X \arrow[r, "g"] & E^\bullet \arrow[r] & U^\bullet[1].
\end{tikzcd}$$

Since $\mathrm{Hom}_{\mathscr{D}^{\mathrm{b}}(A)}((\sigma_{\geqslant 0}E^\bullet)[-1], Y) \simeq \mathrm{Hom}_{\mathscr{K}^{\mathrm{b}}(A)}((\sigma_{\geqslant 0}E^\bullet)[-1], Y) = 0$, the morphism $(w[-1])ra = 0$, and hence there is some morphism $f : X \longrightarrow Y$ in $\mathscr{D}^{\mathrm{b}}(A)$ such that $ra = hf$. Then we have the following commutative diagram in $\mathscr{D}^{\mathrm{b}}(A)$

$$\begin{tikzcd}
& V^\bullet \arrow[dl, Rightarrow, "r"'] \arrow[dr, Rightarrow, "h"] & \\
U^\bullet \arrow[d, Rightarrow, "s"'] \arrow[drr, "a"' near end] & & X \arrow[d, "f"] \arrow[dll, Rightarrow, "1_X"' near start] \\
X & & Y,
\end{tikzcd}$$

which means that the morphisms $X \xLeftarrow{s} U^\bullet \xrightarrow{a} Y$ and $X \xLeftarrow{1_X} X \xrightarrow{f} Y$ in $\mathscr{D}^{\mathrm{b}}(A)/\mathscr{K}^{\mathrm{b}}(\nu_A\text{-Stp})$ are equal. Since the embedding of A-mod into $\mathscr{D}^{\mathrm{b}}(A)$ is fully faithful, the morphism f is given by a morphism in A-mod. Hence the functor Σ' is full, and therefore Σ is a full functor.

Suppose that $f : X \longrightarrow Y$ is a morphism in A-mod such that $\Sigma'(f) = 0$. That is, the morphisms $X \xLeftarrow{1_X} X \xrightarrow{0} Y$ and $X \xLeftarrow{1_X} X \xrightarrow{f} Y$ are equal in $\mathscr{D}^{\mathrm{b}}(A)/\mathscr{K}^{\mathrm{b}}(\nu_A\text{-Stp})$. Then there is a morphism $W^\bullet \xRightarrow{s} X$ such that $sf = 0$ in $\mathscr{D}^{\mathrm{b}}(A)$. Embedding s into a distinguished triangle in $\mathscr{D}^{\mathrm{b}}(A)$, we see that f factorizes in $\mathscr{D}^{\mathrm{b}}(A)$ through a complex in $\mathscr{K}^{\mathrm{b}}(\nu_A\text{-Stp})$, and therefore it follows easily that f factorizes in A-mod through an A-module ν_A-Stp. Hence the functor Σ is faithful.

If A is self-injective, then $\nu_A\text{-Stp} = A\text{-proj}$ and the equivalence was proved by Rickard [14, Theorem 2.1]. If A is not self-injective, then there is a projective A-module P not in ν_A-Stp. Suppose that Σ is an equivalence. Then there is some A-module X such that $X \simeq P[-1]$ in $\mathscr{D}^{\mathrm{b}}(A)/\mathscr{K}^{\mathrm{b}}(\nu_A\text{-Stp})$. That is, there is an isomorphism $X \overset{s}{\Longleftarrow} U^{\bullet} \overset{t}{\Longrightarrow} P[-1]$ in $\mathscr{D}^{\mathrm{b}}(A)/\mathscr{K}^{\mathrm{b}}(\nu_A\text{-Stp})$. Then by Octahedral Axiom, we can form the following commutative diagram in $\mathscr{D}^{\mathrm{b}}(A)$

$$\begin{tikzcd}
E_1^{\bullet} \arrow[r] \arrow[d, equal] & U^{\bullet} \arrow[r, Rightarrow, "t"] \arrow[d, Rightarrow, "s"] & P[-1] \arrow[r] \arrow[d, "h"] & E_1^{\bullet}[1] \arrow[d, equal] \\
E_1^{\bullet} \arrow[r, "g"] & X \arrow[r, "t"] \arrow[d] & \mathrm{con}(g) \arrow[r] \arrow[d] & E_1^{\bullet}[1], \\
 & E_2^{\bullet} \arrow[r, equal] & E_2^{\bullet} &
\end{tikzcd}$$

where $E_1^{\bullet}$ and $E_2^{\bullet}$ are in $\mathscr{K}^{\mathrm{b}}(\nu_A\text{-Stp})$, and $\mathrm{con}(g)$ is the mapping cone of g. From the vertical distinguished triangle on the right side, we see that the mapping cone $\mathrm{con}(h)$ of h is isomorphic in $\mathscr{D}^{\mathrm{b}}(A)$ to a complex $E_2^{\bullet}$ in $\mathscr{K}^{\mathrm{b}}(\nu_A\text{-Stp})$. All the terms of $\mathrm{con}(h)$ in non-zero degrees are in ν_A-Stp and $P \oplus X$ is a direct summand of the 0-degree term of $\mathrm{con}(h)$. Hence P is isomorphic to a complex in $\mathscr{K}^{\mathrm{b}}(\nu_A\text{-Stp})$ which is impossible since P is projective and is not in ν_A-Stp. This finishes the proof. $\square$

Remark: In the above proposition, suppose that P is a projective-injective A-module, if we replace $A\text{-}\underline{\mathrm{mod}}^{\nu}$ by the quotient category of A-mod modulo morphisms factorizing through modules in $\mathrm{add}(P)$, and replace $\mathscr{D}^{\mathrm{b}}(A)/\mathscr{K}^{\mathrm{b}}(\nu_A\text{-Stp})$ by $\mathscr{D}^{\mathrm{b}}(A)/\mathscr{K}^{\mathrm{b}}(\mathrm{add}(P))$, then the proof of Proposition 4.1 actually can be used to show that in this case the functor Σ is also fully faithful.

Now for each iterated almost ν-stable derived equivalence $F : \mathscr{D}^{\mathrm{b}}(A) \longrightarrow \mathscr{D}^{\mathrm{b}}(B)$. By Lemma 3.2, we see that F induces an equivalence between the triangulated categories $\mathscr{D}^{\mathrm{b}}(A)/\mathscr{K}^{\mathrm{b}}(\nu_A\text{-Stp})$ and $\mathscr{D}^{\mathrm{b}}(B)/\mathscr{K}^{\mathrm{b}}(\nu_B\text{-Stp})$. We also denote this equivalence by F. In the following, we will see that there is an equivalence $\phi_F : A\text{-}\underline{\mathrm{mod}}^{\nu} \longrightarrow B\text{-}\underline{\mathrm{mod}}^{\nu}$ such that the diagram

$$\begin{tikzcd}
A\text{-}\underline{\mathrm{mod}}^{\nu} \arrow[r, "\Sigma"] \arrow[d, "\phi_F"] & \mathscr{D}^{\mathrm{b}}(A)/\mathscr{K}^{\mathrm{b}}(\nu_A\text{-Stp}) \arrow[d, "F"] \\
B\text{-}\underline{\mathrm{mod}}^{\nu} \arrow[r, "\Sigma"] & \mathscr{D}^{\mathrm{b}}(B)/\mathscr{K}^{\mathrm{b}}(\nu_B\text{-Stp})
\end{tikzcd}$$

of additive functors is commutative up to isomorphism. Moreover, the functor ϕ_F also induces an equivalence between the stable module categories $A\text{-}\underline{\mathrm{mod}}$ and $B\text{-}\underline{\mathrm{mod}}$.

Before we give the construction of ϕ_F, we give the following lemma, which generalizes [6, Lemma 2.2] and will be used in the construction of ϕ_F.

Lemma 4.2. *Let A be an arbitrary ring, and let A-*Mod *be the category of all left (not necessarily finitely generated) A-modules. Suppose $X^{\bullet}$ is a complex over A-*Mod *bounded above and $Y^{\bullet}$ is a complex over A-*Mod *bounded below. If there is an integer m such that X^i is projective for all $i > m$ and Y^j is injective for all $j < m$, then $\theta_{X^{\bullet},Y^{\bullet}} : \mathrm{Hom}_{\mathscr{K}(A\text{-Mod})}(X^{\bullet},Y^{\bullet}) \to \mathrm{Hom}_{\mathscr{D}(A\text{-Mod})}(X^{\bullet},Y^{\bullet})$ induced by the localization functor $\theta : \mathscr{K}(A\text{-Mod}) \to \mathscr{D}(A\text{-Mod})$ is an isomorphism.*

Proof. Without loss of generality, we can assume that $m = 0$. For simplicity, we write $\mathscr{K}$ for $\mathscr{K}(A\text{-Mod})$ and $\mathscr{D}$ for $\mathscr{D}(A\text{-Mod})$. Also, the Hom-spaces $\mathrm{Hom}_{\mathscr{K}}(-,-)$ and $\mathrm{Hom}_{\mathscr{D}}(-,-)$ will be denoted by ${}_{\mathscr{K}}(-,-)$ and ${}_{\mathscr{D}}(-,-)$, respectively.

First, we show that, for each A-module Z, the induced map

$$\theta_{X^\bullet,Z[1]} : \mathscr{K}(X^\bullet, Z[1]) \longrightarrow \mathscr{D}(X^\bullet, Z[1])$$

is monic. Indeed, applying $\mathscr{K}(-,Z[1])$ and $\mathscr{D}(-,Z[1])$ to the distinguished triangle

$$\sigma_{\geqslant 0}X^\bullet \longrightarrow X^\bullet \longrightarrow \sigma_{<0}X^\bullet \longrightarrow (\sigma_{\geqslant 0}X^\bullet)[1],$$

we get a commutative diagram with exact rows.

$$\begin{tikzcd}
\mathscr{K}(\sigma_{\geqslant 0}X^\bullet, Z) \arrow[r] \arrow[d, "\theta_{\sigma_{\geqslant 0}X^\bullet,Z}"] & \mathscr{K}(\sigma_{<0}X^\bullet, Z[1]) \arrow[r] \arrow[d, "\theta_{\sigma_{<0}X^\bullet,Z[1]}"] & \mathscr{K}(X^\bullet, Z[1]) \arrow[r] \arrow[d, "\theta_{X^\bullet,Z[1]}"] & \mathscr{K}(\sigma_{\geqslant 0}X^\bullet, Z[1]) \arrow[d, "\theta_{\sigma_{\geqslant 0}X^\bullet,Z[1]}"] \\
\mathscr{D}(\sigma_{\geqslant 0}X^\bullet, Z) \arrow[r] & \mathscr{D}(\sigma_{<0}X^\bullet, Z[1]) \arrow[r] & \mathscr{D}(X^\bullet, Z[1]) \arrow[r] & \mathscr{D}(\sigma_{\geqslant 0}X^\bullet, Z[1])
\end{tikzcd}$$

By [6, Lemma 2.2], the maps $\theta_{\sigma_{\geqslant 0}X^\bullet,Z}$ and $\theta_{\sigma_{<0}X^\bullet,Z[1]}$ are isomorphisms. Since $\mathscr{K}(\sigma_{\geqslant 0}X^\bullet, Z[1]) = 0$, the map $\theta_{\sigma_{\geqslant 0}X^\bullet,Z[1]}$ is clearly monic. Thus, by the Five Lemma (see, for example [15, p.13]), the map $\theta_{X^\bullet,Z[1]}$ is monic.

Next, we show that the map

$$\theta_{X^\bullet,(\sigma_{\geqslant 0}Y^\bullet)[1]} : \mathscr{K}(X^\bullet, (\sigma_{\geqslant 0}Y^\bullet)[1]) \longrightarrow \mathscr{D}(X^\bullet, (\sigma_{\geqslant 0}Y^\bullet)[1])$$

is monic. Indeed, applying $\mathscr{K}(X^\bullet,-)$ and $\mathscr{D}(X^\bullet,-)$ to the distinguished triangle

$$Y^0 \longrightarrow (\sigma_{>0}Y^\bullet)[1] \longrightarrow (\sigma_{\geqslant 0}Y^\bullet)[1] \longrightarrow Y^0[1],$$

we get a commutative diagram with exact rows.

$$\begin{tikzcd}
\mathscr{K}(X^\bullet, Y^0) \arrow[r] \arrow[d, "\theta_{\sigma_{X^\bullet,Y^0}}"] & \mathscr{K}(X^\bullet, (\sigma_{>0}Y^\bullet)[1]) \arrow[r] \arrow[d, "\theta_{X^\bullet,(\sigma_{>0}Y^\bullet)[1]}"] & \mathscr{K}(X^\bullet, (\sigma_{\geqslant 0}Y^\bullet)[1]) \arrow[r] \arrow[d, "\theta_{X^\bullet,(\sigma_{\geqslant 0}Y^\bullet)[1]}"] & \mathscr{K}(X^\bullet, Y^0[1]) \arrow[d, "\theta_{X^\bullet,Y^0[1]}"] \\
\mathscr{D}(X^\bullet, Y^0) \arrow[r] & \mathscr{D}(X^\bullet, (\sigma_{<0}Y^\bullet)[1]) \arrow[r] & \mathscr{D}(X^\bullet, (\sigma_{\geqslant 0}Y^\bullet)[1]) \arrow[r] & \mathscr{D}(X^\bullet, Y^0[1])
\end{tikzcd}$$

Again by [6, Lemma 2.2], the left two vertical maps are isomorphisms. By the above discussion, we see that $\theta_{X^\bullet,Y^0[1]}$ is monic. So, by the Five Lemma again, the map $\theta_{X^\bullet,(\sigma_{\geqslant 0}Y^\bullet)[1]}$ is monic.

Finally, applying $\mathscr{K}(X^\bullet,-)$ and $\mathscr{D}(X^\bullet,-)$ to the distinguished triangle

$$(\sigma_{<0}Y^\bullet)[-1] \longrightarrow \sigma_{\geqslant 0}Y^\bullet \longrightarrow Y^\bullet \longrightarrow \sigma_{<0}Y^\bullet,$$

we get a commutative diagram

$$\begin{tikzcd}
\mathscr{K}(X^\bullet, (\sigma_{<0}Y^\bullet)[-1]) \arrow[r] \arrow[d, "\theta_{X^\bullet,(\sigma_{<0}Y^\bullet)[-1]}"] & \mathscr{K}(X^\bullet, \sigma_{\geqslant 0}Y^\bullet) \arrow[r] \arrow[d, "\theta_{X^\bullet,\sigma_{\geqslant 0}Y^\bullet}"] & \mathscr{K}(X^\bullet, Y^\bullet) \arrow[r] \arrow[d, "\theta_{X^\bullet,Y^\bullet}"] & \mathscr{K}(X^\bullet, \sigma_{<0}Y^\bullet) \arrow[r] \arrow[d, "\theta_{X^\bullet,\sigma_{<0}Y^\bullet}"] & \mathscr{K}(X^\bullet, (\sigma_{\geqslant 0}Y^\bullet)[1]) \arrow[d, "\theta_{X^\bullet,(\sigma_{\geqslant 0}Y^\bullet)[1]}"] \\
\mathscr{D}(X^\bullet, (\sigma_{<0}Y^\bullet)[-1]) \arrow[r] & \mathscr{D}(X^\bullet, \sigma_{\geqslant 0}Y^\bullet) \arrow[r] & \mathscr{D}(X^\bullet, Y^\bullet) \arrow[r] & \mathscr{D}(X^\bullet, \sigma_{<0}Y^\bullet) \arrow[r] & \mathscr{D}(X^\bullet, (\sigma_{\geqslant 0}Y^\bullet)[1])
\end{tikzcd}$$

By assumption, the complex $\sigma_{<0}Y^\bullet$ is a bounded complex of injective A-modules. So, the maps $\theta_{X^\bullet,(\sigma_{<0}Y^\bullet)[-1]}$ and $\theta_{X^\bullet,\sigma_{<0}Y^\bullet}$ are isomorphisms. By [6, Lemmma 2.2], the map $\theta_{X^\bullet,\sigma_{\geqslant 0}Y^\bullet}$ is an isomorphism. We have already proved that the map $\theta_{X^\bullet,(\sigma_{\geqslant 0}Y^\bullet)[1]}$ is monic. Then by applying the Five Lemma again, the proof is completed. $\square$

Now we fix some notations for the rest of this section. Let $F : \mathscr{D}^{\mathrm{b}}(A) \longrightarrow \mathscr{D}^{\mathrm{b}}(B)$ be an iterated almost ν-stable derived equivalences between two Artin algebras A and B, and let G be a quasi-inverse of F. Let $T^\bullet$ and $\bar{T}^\bullet$ be the tilting complexes associated to F and G, respectively. Then by Theorem 1.1 and Lemma 3.1, the terms of $T^\bullet$ in non-zero degrees are all in ν_A-Stp, and the terms of $\bar{T}^\bullet$ in non-zero degrees are all in ν_B-Stp.

Keeping these notations, we have the following lemma.

Lemma 4.3. *For each A-module X, the complex $F(X)$ is isomorphic in $\mathscr{D}^{\rm b}(B)$ to a radical complex $\bar{T}_X^\bullet$ with $\bar{T}_X^\pm \in \nu_B$-Stp. Moreover, the complex $\bar{T}_X^\bullet$ of this form is unique up to isomorphism in $\mathscr{C}^b(B)$. In particular, if X is a projective (respectively, injective) module, then $\bar{T}_X^\bullet$ is isomorphic in $\mathscr{C}^b(B)$ to a complex in* $\mathrm{add}(\bar{T}^\bullet)$ *(respectively,* $\mathrm{add}(\nu_B\bar{T}^\bullet)$*).*

Proof. By the proof of Theorem 1.1, we see that $F \simeq F_2F_1^{-1}$ for two almost ν-stable derived equivalences $F_1 : \mathscr{D}^{\rm b}(C) \longrightarrow \mathscr{D}^{\rm b}(A)$ and $F_2 : \mathscr{D}^{\rm b}(C) \longrightarrow \mathscr{D}^{\rm b}(B)$. For each A-module X, by [6, Lemma 3.2] and the definition of almost ν-stable derived equivalences, we see that $F_1^{-1}(X)$ is isomorphic in $\mathscr{D}^{\rm b}(C)$ to a complex $Q_X^\bullet$ with $Q_X^i = 0$ for all $i > 0$ and $Q_X^i \in \nu_C$-Stp for all $i < 0$. Applying F_2 to the distinguished triangle $(\sigma_{<0}Q_X^\bullet)[-1] \longrightarrow Q_X^0 \longrightarrow Q_X^\bullet \longrightarrow \sigma_{<0}Q_X^\bullet$, we get a distinguished triangle in $\mathscr{D}^{\rm b}(B)$

$$F_2(\sigma_{<0}Q_X^\bullet)[-1] \longrightarrow F_2(Q_X^0) \longrightarrow F_2(Q_X^\bullet) \longrightarrow F_2(\sigma_{<0}Q_X^\bullet).$$

Since $(\sigma_{<0}Q_X^\bullet)[-1]$ is a complex in $\mathscr{K}^{\rm b}(\nu_C\text{-Stp})$, by Lemma 3.2, the complex $F_2(\sigma_{<0}Q_X^\bullet)[-1]$ is isomorphic in $\mathscr{D}^{\rm b}(B)$ to a complex $U^\bullet$ in $\mathscr{K}^{\rm b}(\nu_B\text{-Stp})$. By [6, Lemma 3.1] and the definition of almost ν-stable derived equivalences, the complex $F_2(Q_X^0)$ is isomorphic in $\mathscr{D}^{\rm b}(B)$ to a complex $V^\bullet$ with $V^i \in \nu_B$-Stp for all $i > 0$ and $V^i = 0$ for all $i < 0$. Thus, the complex $F_2(Q_X^\bullet)$, which is isomorphic in $\mathscr{D}^{\rm b}(B)$ to $F(X)$, is isomorphic in $\mathscr{D}^{\rm b}(B)$ to the mapping cone $\mathrm{con}(\alpha)$ of a chain map α from $U^\bullet$ to $V^\bullet$. Now it is clear that all the terms of $\mathrm{con}(\alpha)$ in non-zero degrees are in ν_B-Stp. Taking a radical complex $\bar{T}_X^\bullet$ which is isomorphic to $\mathrm{con}(\alpha)$ in $\mathscr{K}^{\rm b}(B)$, we see that $F(X)$ is isomorphic to $\bar{T}_X^\bullet$ and $\bar{T}_X^\pm \in \nu_B$-Stp.

Suppose that $W^\bullet$ is another radical complex with $W^\pm \in \nu_B$-Stp, and $F(X) \simeq W^\bullet$. Then $W^\bullet$ and $\bar{T}_X^\bullet$ are isomorphic in $\mathscr{D}^{\rm b}(B)$. By Lemma 4.2, they are isomorphic in $\mathscr{K}^{\rm b}(B)$. Since both $W^\bullet$ and $\bar{T}_X^\bullet$ are radical complexes, they are also isomorphic in $\mathscr{C}^b(B)$.

Since all the complexes in $\mathrm{add}(\bar{T}^\bullet)$ and $\mathrm{add}(\nu_B\bar{T}^\bullet)$ have the desired form, the last statement follows by the uniqueness of $\bar{T}_X^\bullet$. $\square$

In the following, without loss of generality, we fix for each A-module X a complex $\bar{T}_X^\bullet$ defined in Lemma 4.3 and assume that $F(X) = \bar{T}_X^\bullet$ for all A-modules X. Let X and Y be two A-modules. There is a natural isomorphism

$$\mathrm{Hom}_A(X,Y) \simeq \mathrm{Hom}_{\mathscr{D}^{\rm b}(B)}(\bar{T}_X^\bullet, \bar{T}_Y^\bullet)$$

sending f to $F(f)$. By Lemma 4.2, there is a natural isomorphism

$$\mathrm{Hom}_{\mathscr{K}^{\rm b}(B)}(\bar{T}_X^\bullet, \bar{T}_Y^\bullet) \simeq \mathrm{Hom}_{\mathscr{D}^{\rm b}(B)}(\bar{T}_X^\bullet, \bar{T}_Y^\bullet)$$

induced by the localization functor from $\mathscr{K}^{\rm b}(B)$ to $\mathscr{D}^{\rm b}(B)$. It is easy to see that there is a natural map

$$\mathrm{Hom}_{\mathscr{K}^{\rm b}(B)}(\bar{T}_X^\bullet, \bar{T}_Y^\bullet) \longrightarrow \underline{\mathrm{Hom}}_B^\nu(\bar{T}_X^0, \bar{T}_Y^0)$$

sending $u^\bullet$ to $\underline{u}^0$. Indeed, if $u^\bullet = v^\bullet$ in $\mathrm{Hom}_{\mathscr{K}^{\rm b}(B)}(\bar{T}_X^\bullet, \bar{T}_Y^\bullet)$, then $u^0 - v^0$ factorizes through $\bar{T}_X^1 \oplus \bar{T}_Y^{-1}$ which is in ν_B-Stp by definition. This means $\underline{u}^0 - \underline{v}^0 = 0$ in $\underline{\mathrm{Hom}}_B^\nu(\bar{T}_X^0, \bar{T}_Y^0)$. Altogether, we have a natural morphism

$$\phi : \mathrm{Hom}_A(X,Y) \longrightarrow \underline{\mathrm{Hom}}_B^\nu(\bar{T}_X^0, \bar{T}_Y^0)$$

sending f to $\underline{u}^0$, where $u^\bullet$ is a chain map such that $u^\bullet = F(f)$. Now if f factorizes through an A-module in ν_A-Stp, then $u^\bullet$ factorizes through a complex $P^\bullet$ in $\mathscr{K}^{\rm b}(\nu_B\text{-Stp})$ by Lemma 3.2. By Lemma 4.2, we can assume that $u^\bullet = g^\bullet h^\bullet$ in $\mathscr{K}^{\rm b}(B)$ for chain maps $g^\bullet : \bar{T}_X^\bullet \longrightarrow P^\bullet$ and $h^\bullet : P^\bullet \longrightarrow \bar{T}_Y^\bullet$. Thus, it follows that $u^0 - g^0h^0$ factorizes through $\bar{T}_X^1 \oplus \bar{T}_Y^{-1}$, and consequently u^0 factorizes through $P^0 \oplus \bar{T}_X^1 \oplus \bar{T}_Y^{-1}$ which is in ν_B-Stp. Hence $\underline{u}^0 = 0$. Hence we get a natural morphism

$$\bar{\phi} : \underline{\mathrm{Hom}}_A^\nu(X,Y) \longrightarrow \underline{\mathrm{Hom}}_B^\nu(\bar{T}_X^0, \bar{T}_Y^0)$$

Now we define a functor $\phi_F : A\text{-}\underline{\mathrm{mod}}^{\nu} \longrightarrow B\text{-}\underline{\mathrm{mod}}^{\nu}$. For each A-module X, we set $\phi_F(X) := \bar{T}_X^0$, and for each morphism $\underline{f} \in \underline{\mathrm{Hom}}_A^{\nu}(X,Y)$, we define $\phi_F(\underline{f}) := \underline{u}^0$, where $u^\bullet = F(f)$. Now it is easy to see that the diagram

$$\begin{tikzcd}
A\text{-}\underline{\mathrm{mod}}^{\nu} \arrow[r, "\Sigma"] \arrow[d, "\phi_F"] & \mathscr{D}^{\mathrm{b}}(A)/\mathscr{K}^{\mathrm{b}}(\nu_A\text{-Stp}) \arrow[d, "F"] & (\clubsuit) \\
B\text{-}\underline{\mathrm{mod}}^{\nu} \arrow[r, "\Sigma"] & \mathscr{D}^{\mathrm{b}}(B)/\mathscr{K}^{\mathrm{b}}(\nu_B\text{-Stp}) &
\end{tikzcd}$$

is commutative up to isomorphism. Indeed, one can check that the isomorphism $\bar{T}_X^\bullet \overset{s}{\Longleftarrow} \sigma_{\geqslant 0}\bar{T}_X^\bullet \overset{t}{\Longrightarrow} X$ in $\mathscr{D}^{\mathrm{b}}(B)/\mathscr{K}^{\mathrm{b}}(\nu_B\text{-Stp})$ with s and t the canonical maps is a natural map, and this gives rise to an isomorphism from the functor $F\Sigma$ to the functor $\Sigma\phi_F$.

For an Artin algebra, in the following theorem, we denote by $A\text{-}\overline{\mathrm{mod}}$ the quotient category of A-mod modulo morphisms factorizing through injective modules.

Theorem 4.4. *Let $F : \mathscr{D}^{\mathrm{b}}(A) \longrightarrow \mathscr{D}^{\mathrm{b}}(B)$ be an iterated almost ν-stable derived equivalence. Then we have the following:*

(1) *The functor $\phi_F : A\text{-}\underline{\mathrm{mod}}^{\nu} \longrightarrow B\text{-}\underline{\mathrm{mod}}^{\nu}$ is an equivalence;*

(2) *The functor ϕ_F induces an equivalence between $A\text{-}\overline{\mathrm{mod}}$ and $B\text{-}\overline{\mathrm{mod}}$;*

(3) *The functor ϕ_F induces an equivalence between $A\text{-}\underline{\mathrm{mod}}$ and $B\text{-}\underline{\mathrm{mod}}$;*

(4) *The functor ϕ_F is uniquely (up to isomorphism) determined by the commutative diagram $(\clubsuit)$. Moreover, if $F' : \mathscr{D}^{\mathrm{b}}(B) \longrightarrow \mathscr{D}^{\mathrm{b}}(C)$ is another iterated almost ν-stable derived equivalence, then $\phi_{F'F} \simeq \phi_{F'}\phi_F$.*

Proof. Let G be a quasi-inverse of F. Then G also induces an equivalence between $\mathscr{D}^{\mathrm{b}}(B)/\mathscr{K}^{\mathrm{b}}(\nu_B\text{-Stp})$ and $\mathscr{D}^{\mathrm{b}}(A)/\mathscr{K}^{\mathrm{b}}(\nu_A\text{-Stp})$. We also denote it by G. Then by the above commutative diagram of additive functors, the functor $\Sigma\phi_G\phi_F$ is isomorphic to the functor $GF\Sigma$, which is isomorphic to Σ. By Proposition 4.1, the functor Σ is a fully faithful embedding. Hence $\phi_G\phi_F$ is isomorphic to $1_{A\text{-}\underline{\mathrm{mod}}^{\nu}}$. By symmetry, the functor $\phi_F\phi_G$ is also isomorphic to $1_{B\text{-}\underline{\mathrm{mod}}^{\nu}}$. Hence ϕ_F is an equivalence, and (1) is proved.

By the construction of ϕ_F, it follows from Lemma 4.3 that ϕ_F sends projective modules to projective modules, and sends injective modules to injective modules. Moreover, the modules in ν_A-Stp and ν_B-Stp are all projective-injective. Thus, the statements (2) and (3) follow.

(4) If $\phi : A\text{-}\underline{\mathrm{mod}}^{\nu} \longrightarrow B\text{-}\underline{\mathrm{mod}}^{\nu}$ is a functor such that $\Sigma\phi \simeq F\Sigma$, then the functor $\Sigma\phi$ is isomorphic to $\Sigma\phi_F$. Hence $\phi \simeq \phi_F$ since Σ is fully faithful. The rest of (4) follows similarly. ☐

Remark: (1) It follows from the definition of iterated almost ν-stable derived equivalences and [6, Theorem 3.7] that every iterated almost ν-stable derived equivalence induces an equivalence between the stable module categories, however, the proof of Theorem 4.4 presented here is not based on the earlier result [6, Thoerem 3.7], and is completely different from the proof there. Moreover, Theorem 4.4 is more general than [6, Theorem 3.7] since we get an equivalence between $A\text{-}\underline{\mathrm{mod}}^{\nu}$ and $B\text{-}\underline{\mathrm{mod}}^{\nu}$ which is not obtained in [6, Theorem 3.7].

(2) In case that F is an almost ν-stable derived equivalence, it follows by definition that the stable equivalence from $A\text{-}\underline{\mathrm{mod}}$ to $B\text{-}\underline{\mathrm{mod}}$ induced by the functor ϕ_F coincides with the stable functor $\bar{F}$ considered in [10].

5 Constructions of iterated almost ν-stable derived equivalences

In this section, we shall give some constructions of iterated almost ν-stable derived equivalences.

Let us recall from [2] the definition of approximations. Let $\mathcal{C}$ be a category, and let $\mathcal{D}$ be a full subcategory of $\mathcal{C}$, and X an object in $\mathcal{C}$. A morphism $f : D \longrightarrow X$ in $\mathcal{C}$ is called a *right $\mathcal{D}$-approximation* of X if $D \in \mathcal{D}$ and the induced map $\mathrm{Hom}_{\mathcal{C}}(D', f)$: $\mathrm{Hom}_{\mathcal{C}}(D',D) \longrightarrow \mathrm{Hom}_{\mathcal{C}}(D',X)$ is surjective for every object $D' \in \mathcal{D}$. Dually, one can define *left $\mathcal{D}$-approximations*.

By Theorem 1.1, to get an iterated almost ν-stable derived equivalence, we only need to construct a derived equivalence with the involved tilting complexes satisfying the conditions in Theorem 1.1. Let A be an algebra, and let P, Q be two projective A-modules satisfying the following two conditions:

(1) $\mathrm{add}({}_AP) = \mathrm{add}(\nu_AP)$, $\mathrm{add}({}_AQ) = \mathrm{add}(\nu_AQ)$;

(2) $\mathrm{Hom}_A(P, Q) = 0$.

For each positive integer r, we can form the following complex:

$$0 \longrightarrow P^{-r} \xrightarrow{f_r}\longrightarrow P^{-r+1} \longrightarrow \cdots \longrightarrow P^{-1} \xrightarrow{f_1} A \longrightarrow 0,$$

where $f_1 : P^{-1} \longrightarrow A$ is a right $\mathrm{add}({}_AP)$-approximation of A, and $f_{i+1} : P^{-i-1} \longrightarrow P^{-i}$ is a right $\mathrm{add}({}_AP)$-approximation of $\mathrm{Ker}(f_i)$ for $i = 1, \cdots, r-1$. Similarly, we can form a complex

$$0 \longrightarrow A \xrightarrow{g_1} Q^1 \longrightarrow \cdots \longrightarrow Q^{s-1} \xrightarrow{g_s} Q^s \longrightarrow 0,$$

where g_1 is a left $\mathrm{add}({}_AQ)$-approximation of A, and g_{i+1} is a left $\mathrm{add}({}_AQ)$-approximation of $\mathrm{Coker}(g_i)$ for $i = 1, 2, \cdots, s-1$. Since $\mathrm{Hom}_A(P, Q) = 0$, connecting the two complexes together, we get a complex

$$0 \longrightarrow P^{-r} \longrightarrow \cdots \longrightarrow P^{-1} \xrightarrow{f_1} A \xrightarrow{g_1} Q^1 \longrightarrow \cdots \longrightarrow Q^s \longrightarrow 0,$$

where A is in degree zero. We denote this complex by $T^\bullet_{P,Q}$, and let $T^\bullet := T^\bullet_{P,Q} \oplus P[r] \oplus Q[-s]$.

Proposition 5.1. *Keeping the notations above, we have the following:*

(1) *The complex $T^\bullet$ is a tilting complex.*

(2) *Let $B := \mathrm{End}_{\mathscr{D}^{\mathrm{b}}(A)}(T^\bullet)$. Then $T^\bullet$ induces an iterated almost ν-stable derived equivalence between the algebras A and B.*

Proof. (1) By the construction of $T^\bullet$, we have

$$T^i = \begin{cases} P^{-r} \oplus P, & i = -r; \\ P^i, & -r < i < 0; \\ A, & i = 0; \\ Q^i, & 0 < i < s; \\ Q^s \oplus Q, & i = s; \\ 0 & \text{otherwise.} \end{cases}, \text{ and } d_T^i = \begin{cases} \left[\begin{smallmatrix} f_r \\ 0 \end{smallmatrix}\right], & i = -r; \\ f_{-i}, & -r < i < 0; \\ g_{i+1}, & 0 \leqslant i < s-1; \\ \left[\begin{smallmatrix} g_s \\ 0 \end{smallmatrix}\right], & i = s-1; \\ 0 & \text{otherwise.} \end{cases}$$

We first show that $\mathrm{Hom}_{\mathscr{K}^{\mathrm{b}}(A\text{-proj})}(T^\bullet, T^\bullet[i]) = 0$ for all $i \neq 0$. Assume that i is a positive integer. Let $u^\bullet$ be a morphism in $\mathrm{Hom}_{\mathscr{K}^{\mathrm{b}}(A\text{-proj})}(T^\bullet, T^\bullet[i])$. Then we have the following commutative diagram

$$\begin{tikzcd}
\cdots \arrow[r] & T^{-i-1} \arrow[r, "d_T^{-i-1}"] \arrow[d, "u^{-i-1}"] & T^{-i} \arrow[r, "d_T^{-i}"] \arrow[d, "u^{-i}"] & T^{-i+1} \arrow[r, "d_T^{-i+1}"] \arrow[d, "u^{-i+1}"] & \cdots \arrow[r] & T^{-1} \arrow[r, "d^{-1}"] \arrow[d, "u^{-1}"] & T^0 \arrow[r, "d_T^0"] \arrow[d, "u^0"] & T^1 \arrow[r, "d_T^1"] \arrow[d, "u^1"] & \cdots \\
\cdots \arrow[r] & T^{-1} \arrow[r, "d_T^{-1}"] & T^0 \arrow[r, "d_T^0"] & T^1 \arrow[r, "d_T^1"] & \cdots \arrow[r] & T^{i-1} \arrow[r, "d_T^{i-1}"] & T^i \arrow[r, "d_T^i"] & T^{i+1} \arrow[r, "d_T^{i+1}"] & \cdots
\end{tikzcd}$$

Since $\mathrm{Hom}_A(P, Q) = 0$, we have $u^k = 0$ for all $-i < k < 0$. By definition, $T^{-i} \in \mathrm{add}({}_AP)$. Since $d_T^{-1} = f_1$ is a right $\mathrm{add}({}_AP)$-approximation, there is a map $h^{-i} : T^{-i} \longrightarrow T^{-1}$ such that $u^{-i} = h^{-i}d_T^{-1}$. Thus, $(u^{-i-1} - d_T^{-i-1}h^{-i})d_T^{-1} = d_T^{-i-1}u^{-i} - d_T^{-i-1}h^{-i}d_T^{-1} = d_T^{-i-1}u^{-i} - d_T^{-i-1}u^{-i} = 0$. Since d_T^{-2} is a right $add({}_AP)$-approximation of $\mathrm{Ker}(d_T^{-1})$, there is a map $h^{-i-1} : T^{-i-1} \longrightarrow T^{-2}$ such that $u^{-i-1} - d_T^{-i-1}h^{-i} = h^{-i-1}d_T^{-2}$, that is $u^{-i-1} = d_T^{-i-1}h^{-i} + h^{-i-1}d_T^{-2}$. Similarly, for each integer $k < -i-1$, there are maps $h^{k+1} : T^{k+1} \longrightarrow T^{k+i}$ and $h^k : T^k \longrightarrow T^{k+i-1}$ such that $u^k = d_T^k h^{k+1} + h^k d_T^{k+i-1}$. Defining $h^k = 0$ for all $-i < k \leqslant 0$, we have $u^k = d_T^k h^{k+1} + h^k d_T^{k+i-1}$

for all $k < 0$. Similarly, we can prove that $u^k = d_T^k h^{k+1} + h^k d_T^{k+i-1}$ for $k \geqslant 0$. Altogether, we have shown that $u^\bullet = 0$ in $\mathscr{K}^{\mathrm{b}}(A\text{-proj})$. Hence $\mathrm{Hom}_{\mathscr{K}^{\mathrm{b}}(A\text{-proj})}(T^\bullet, T^\bullet[i]) = 0$ for all $i > 0$. By an analogous proof, we have $\mathrm{Hom}_{\mathscr{K}^{\mathrm{b}}(A\text{-proj})}(T^\bullet, T^\bullet[i]) = 0$ for all $i < 0$. Finally, since $P[r]$ and $Q[-s]$ are in $\mathrm{add}(T^\bullet)$, we deduce that ${}_AA$ is in the triangulated subcategory of $\mathscr{K}^{\mathrm{b}}(A\text{-proj})$ generated by $\mathrm{add}(T^\bullet)$. Hence $\mathrm{add}(T^\bullet)$ generates $\mathscr{K}^{\mathrm{b}}(A\text{-proj})$ as a triangulated category, and consequently $T^\bullet$ is a tilting complex over A.

(2) Let $F : \mathscr{D}^{\mathrm{b}}(A) \longrightarrow \mathscr{D}^{\mathrm{b}}(B)$ be a derived equivalence induced by $T^\bullet$. Also, we use F to denote the equivalence between $\mathscr{K}^{\mathrm{b}}(A\text{-proj})$ and $\mathscr{K}^{\mathrm{b}}(B\text{-proj})$ induced by F. Set $\bar{P} := \mathrm{Hom}_{\mathscr{K}^{\mathrm{b}}(A\text{-proj})}(T^\bullet, P[r]) \simeq F(P[r])$, $\bar{Q} := \mathrm{Hom}_{\mathscr{K}^{\mathrm{b}}(A\text{-proj})}(T^\bullet, Q[-s]) \simeq F(Q[-s])$, and $\bar{U} := \mathrm{Hom}_{\mathscr{K}^{\mathrm{b}}(A\text{-proj})}(T^\bullet, T^\bullet_{P,Q}) \simeq F(T^\bullet_{P,Q})$. Then $F(P) \simeq \bar{P}[-r]$ and $F(Q) \simeq \bar{Q}[s]$. For simplicity, we list some subcomplexes of $T^\bullet$:

$$
\begin{aligned}
P^\bullet: &\quad 0 \longrightarrow P^{-r} \longrightarrow \cdots \longrightarrow P^{-1} \longrightarrow 0,\\
Q^\bullet: &\quad 0 \longrightarrow Q^{1} \longrightarrow \cdots \longrightarrow Q^{s} \longrightarrow 0,\\
R^\bullet: &\quad 0 \longrightarrow A \longrightarrow Q^{1} \longrightarrow \cdots \longrightarrow Q^{s} \longrightarrow 0.
\end{aligned}
$$

By Lemma 2.1, $F(P^\bullet)$ is isomorphic to a complex $\bar{P}^\bullet$ in $\mathscr{K}^{\mathrm{b}}(\mathrm{add}(\bar{P}))$ such that $\bar{P}^i = 0$ for all $i < 0$ and all $i \geqslant r$. The complex $F(Q^\bullet)$ is isomorphic to a complex $\bar{Q}^\bullet$ in $\mathscr{K}^{\mathrm{b}}(\mathrm{add}(\bar{Q}))$ with $\bar{Q}^i = 0$ for all $i > 0$ and all $i \leqslant -s$. Note that there is a distinguished triangle in $\mathscr{K}^{\mathrm{b}}(A\text{-proj})$

$$P^\bullet[-1] \longrightarrow R^\bullet \longrightarrow T^\bullet_{P,Q} \longrightarrow P^\bullet.$$

Applying F, we get a distinguished triangle in $\mathscr{K}^{\mathrm{b}}(B\text{-proj})$:

$$F(P^\bullet)[-1] \longrightarrow F(R^\bullet) \longrightarrow F(T^\bullet_{P,Q}) \longrightarrow F(P^\bullet).$$

Hence $F(R^\bullet)$ is isomorphic to a complex of the following form:

$$0 \longrightarrow \bar{U} \longrightarrow \bar{P}^0 \longrightarrow \bar{P}^1 \longrightarrow \cdots$$

with $\bar{U}$ in degree 0. Next we have a distinguished triangle

$$Q^\bullet \longrightarrow R^\bullet \longrightarrow A \longrightarrow Q^\bullet[1]$$

in $\mathscr{K}^{\mathrm{b}}(A\text{-proj})$. Applying F, we see that $F(A)$ is isomorphic to a complex $\bar{T}^\bullet$ of the form

$$\cdots \longrightarrow \bar{Q}^{-1} \longrightarrow \bar{Q}^0 \longrightarrow \bar{U} \longrightarrow \bar{P}^0 \longrightarrow \bar{P}^1 \longrightarrow \cdots,$$

where $\bar{U}$ is in degree zero. Note that $\bar{T}^\bullet$ is a tilting complex associated to F^{-1} since $F^{-1}(\bar{T}^\bullet) \simeq A$. Since $\mathrm{add}({}_AP) = \mathrm{add}(\nu_A P)$ and $\mathrm{add}({}_AQ) = \mathrm{add}(\nu_A Q)$, we have $\mathrm{add}({}_B\bar{P}) = \mathrm{add}(\nu_B \bar{P})$ and $\mathrm{add}({}_B\bar{Q}) = \mathrm{add}(\nu_B \bar{Q})$. Thus, we have $\mathrm{add}({}_AT^\pm) = \mathrm{add}(\nu_A T^\pm)$ and $\mathrm{add}({}_B\bar{T}^\pm) = \mathrm{add}(\nu_B \bar{T}^\pm)$. By Theorem 1.1, the statement (2) follows. $\square$

To illustrate Proposition 5.1, we give an example. Let A be the finite-dimensional k-algebra given by the quiver

$$\begin{tikzcd}
\underset{1}{\bullet} \arrow[r, shift left, "\alpha"] & \underset{2}{\bullet} \arrow[l, shift left, "\alpha'"] \arrow[r, shift left, "\beta"] & \underset{3}{\bullet} \arrow[l, shift left, "\beta'"] \arrow[r, shift left, "\gamma"] & \underset{4}{\bullet} \arrow[l, shift left, "\gamma'"]
\end{tikzcd}$$

with relations $\alpha'\alpha = \beta\beta' = \alpha\beta = \beta\gamma = \beta'\alpha' = \gamma'\beta' = \beta'\beta - \gamma\gamma' = 0$. We use P_i to denote the indecomposable projective A-module corresponding to the vertex i for $i = 1, 2, 3, 4$. The Loewy structure of the projective A-modules can be listed as follows.

$$
P_1: \begin{array}{c} 1 \\ 2 \\ 1 \end{array} \qquad
P_2: \begin{array}{ccc} & 2 & \\ 1 & & 3 \\ & & \end{array} \qquad
P_3: \begin{array}{ccc} & 3 & \\ 2 & & 4 \\ & 3 & \end{array} \qquad
P_4: \begin{array}{c} 4 \\ 3 \\ 4 \end{array}
$$

Let $P := P_1$ and $Q := P_3 \oplus P_4$. Then we have $\mathrm{add}({}_AP) = \mathrm{add}(\nu_A P)$, $\mathrm{add}({}_AQ) = \mathrm{add}(\nu_A Q)$, and $\mathrm{Hom}_A(P,Q) = 0$. Using Proposition 5.1, we have a tilting complex $T^\bullet$ over A. The indecomposable direct summands of $T^\bullet$ are:

$$\begin{array}{lllllllll} T_1^\bullet: & 0 & \longrightarrow & P_1 & \longrightarrow & 0 & & & \\ T_2^\bullet: & 0 & \longrightarrow & P_1 & \longrightarrow & P_2 & \longrightarrow & P_3 & \longrightarrow 0 \\ T_3^\bullet: & & & & & 0 & \longrightarrow & P_3 & \longrightarrow 0 \\ T_4^\bullet: & & & & & 0 & \longrightarrow & P_4 & \longrightarrow 0 \end{array}$$

A calculation shows that the algebra $B := \mathrm{End}_{\mathscr{D}^{\mathrm{b}}(A)}(T^\bullet)$ is given by the quiver



with relations $\alpha'\alpha = \alpha\beta = \delta\alpha' = \beta\gamma\delta = \gamma\delta\beta\gamma = 0$. By Proposition 5.1, $T^\bullet$ induces an iterated almost ν-stable derived equivalence between A and B. Therefore, A and B are also stably equivalent of Morita type.

The following proposition shows how we can construct iterated almost ν-stable derived equivalences inductively.

Proposition 5.2. *Let $F : \mathscr{D}^{\mathrm{b}}(A) \longrightarrow \mathscr{D}^{\mathrm{b}}(B)$ be an iterated almost ν-stable derived equivalence between two finite-dimensional algebras A and B over a field k, and let ϕ_F be the stable equivalence induced by F (see,* Theorem 4.4*). Then we have the following:*

(1) *For each A-module X, there is an iterated almost ν-stable derived equivalence between the endomorphism algebras $\mathrm{End}_A(A \oplus X)$ and $\mathrm{End}_B(B \oplus \phi_F(X))$;*

(2) *For a finite-dimension self-injective k-algebra C, there is an iterated almost ν-stable derived equivalence between $A \otimes_k C$ and $B \otimes_k C$.*

Proof. Suppose that $F \simeq F_1F_2\cdots F_n$ such that F_i or F_i^{-1} is almost ν-stable for all i. By Theorem 4.4, we have $\phi_F \simeq \phi_{F_1}\phi_{F_2}\cdots\phi_{F_n}$. By the remark after Theorem 4.4, we know that ϕ_{F_i} coincides with the $\overline{F_i}$ considered in [6] for all i. Thus, the statements (1) follows from [6, Corollary 1.3]. The proof of (2) is similar to that of [6, Proposition 6.2]. □

Let us recall from [7] the definition of Φ-Auslander-Yoneda algebras. A subset Φ of the set of natural numbers $\mathbb{N}$ is called admissible provided that: (1) $0 \in \Phi$; (2) If $i+j+k \in \Phi$ for $i,j,k \in \Phi$, then $i+j \in \Phi$ implies that $j+k \in \Phi$. For instance, the sets $\mathbb{N}$, $\{0,1,\cdots,n\}$ are admissible subsets of $\mathbb{N}$. Suppose that Φ be an admissible subset of $\mathbb{N}$. Let A be an Artin algebra, and let X be an A-module. Now we consider the Yoneda algebra $\mathrm{Ext}_A^*(X,X) = \bigoplus_{i\geqslant 0}\mathrm{Hom}_{\mathscr{D}^{\mathrm{b}}(A)}(X,X[i])$ of X, and define $\mathrm{E}_A^\Phi(X) := \bigoplus_{i\in\Phi}\mathrm{Hom}_{\mathscr{D}^{\mathrm{b}}(A)}(X,X[i])$ with multiplication: for $a_i \in \mathrm{Hom}_{\mathscr{D}^{\mathrm{b}}(A)}(X,X[i])$ and $a_j \in \mathrm{Hom}_{\mathscr{D}^{\mathrm{b}}(A)}(X,X[j])$, we define $a_i \cdot a_j = a_ia_j$ if $i+j \in \Phi$, and zero otherwise. Then one can check that $\mathrm{E}_A^\Phi(X)$ is an associated algebra. If $\Phi = \{0\}$, then $\mathrm{E}_A^\Phi(X)$ is isomorphic to $\mathrm{End}_A(X)$. If $\Phi = \mathbb{N}$, then $\mathrm{E}_A^\Phi(X)$ is just the Yoneda algebra of X.

Proposition 5.3. *Let $F : \mathscr{D}^{\mathrm{b}}(A) \longrightarrow \mathscr{D}^{\mathrm{b}}(B)$ be an iterated almost ν-stable derived equivalence between two Artin algebras A and B. Suppose that Φ is an admissible subset of $\mathbb{N}$. Then we have the following:*

(1) *For any A-module X, there is a derived equivalence between the Φ-Auslander-Yoneda algebras $\mathrm{E}_A^\Phi(A \oplus X)$ and $\mathrm{E}_B^\Phi(B \oplus \phi_F(X))$;*

(2) *If Φ is a finite set, then for any A-module X, there is an iterated almost ν-stable derived equivalence between $\mathrm{E}_A^\Phi(A \oplus X)$ and $\mathrm{E}_B^\Phi(B \oplus \phi_F(X))$.*

Proof. Using the result [7, Theorem 3.4], the proof is similar to that of Proposition 5.2 (1). □

Acknowledgement

This work is partially supported by China Postdoctoral Science Foundation (No. 20080440003). The author also thanks the Alexander von Humboldt Foundation for support.

References

[1] S. AL-NOFAYEE and J. RICKARD, Rigidity of tilting complexes and derived equivalence for self-injective algebras. preprint, 2004.

[2] M. AUSLANDER and S. O. SMALØ, Preprojective modules over Artin algebras. *J. Algebra* **66** (1980) 61-122.

[3] M. BROUÉ, Equivalences of blocks of group algebras. In: "*Finite dimensional algebras and related topics*", edited by V. Dlab and L. L. Scott, Kluwer, 1994, 1-26.

[4] D. HAPPEL, *Triangulated categories in the representation theory of finite dimensional algebras.* Cambridge Univ. Press, Cambridge, 1988.

[5] M. HOSHINO and Y. KATO, Tilting complexes associated with a sequence of idempotents. J. Algebra 183(2003), 105-124.

[6] W. HU and C. C. XI, Derived equivalences and stable equivalences of Morita type, I. *Nagoya Math. J.* to appear. Available at: http://arxiv.org/abs/0810.4761

[7] W. HU and C. C. XI, Derived equivalences for Φ-Auslander-Yoneda algebras. preprint 1-27, 2009. Available at: http://arxiv.org/abs/0912.0647

[8] Y. M. LIU and C. C. XI, Constructions of stable equivalences of Morita type for finite dimensional algebras, I. *Trans. Amer. Math. Soc.* **358**(2006), no.6, 2537-2560.

[9] Y. M. LIU and C. C. XI, Constructions of stable equivalences of Morita type for finite dimensional algebras, II. *Math. Z.* **251**(2005) 21-39.

[10] Y. M. LIU and C. C. XI, Constructions of stable equivalences of Morita type for finite dimensional algebras, III. *J. London Math. Soc.* **76** (2007) 567-585.

[11] A. NEEMAN, *Triangulated categories*. Princeton University Press, Princeton and Oxford, 2001.

[12] J. RICKARD, Morita theory for derived categories. *J. London Math. Soc.* **39**(1989) 436-456.

[13] J. RICKARD, Derived equivalence as derived functors. *J. London Math. Soc.* **43**(1991) 37-48.

[14] J. RICKARD, Derived categories and stable equivalences. *J. Pure Appl. Algebra* **64**(1989) 303-317.

[15] C. A. WEIBEL, *An introduction to homological algebra.* Cambridge Univ. Press, 1994.

WEI HU,
School of Mathematical Sciences, Beijing Normal University, Beijing, 100875, China